



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 211/26, 211/70, A61K 31/445	A1	(11) International Publication Number: WO 95/08535 (43) International Publication Date: 30 March 1995 (30.03.95)
(21) International Application Number: PCT/EP94/03121 (22) International Filing Date: 16 September 1994 (16.09.94) (30) Priority Data: 9319534.5 22 September 1993 (22.09.93) GB (71) Applicant (for all designated States except US): THE BOOTS COMPANY PLC [GB/GB]; 1 Thane Road West, Nottingham NG2 3AA (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): RAFFERTY, Paul [GB/GB]; The Boots Company plc, 1 Thane Road West, Nottingham NG2 3AA (GB). TOMETZKI, Gerald, Bernard [GB/GB]; The Boots Company plc, 1 Thane Road West, Nottingham NG2 3AA (GB). (74) Agent: MILLER, Thomas, Kerr; The Boots Company plc, Patents Dept., R4 Pennyfoot Street, Nottingham NG2 3AA (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: 4-ALKYLAMINO-1-PHENOXYALKYL PIPERIDINES AS ANTIINFLAMMATORY, ANTIALLERGIC AND IMMUNOMODULATING AGENTS <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div> (57) Abstract <p>Compounds of formula (I) and pharmaceutically acceptable salts thereof in which R₁, R₂ and R₃ independently represent hydrogen, hydroxy, halo, alkyl or alkoxy; ALK¹ represents a C₂₋₆ alkylene chain optionally substituted by one or more C₁₋₂ alkyl groups; Y represents a piperidine ring which is attached through nitrogen to ALK¹; R₄ represents hydrogen or a C₁₋₄ alkyl group; the broken line in --- represents a bond, or is absent, and the free valency on Y is taken up by hydrogen and the free valency on CR₄ is taken up by hydrogen or a C₁₋₄ alkyl group; ALK² is absent or represents a C₁₋₄ alkylene chain optionally substituted by one or more C₁₋₂ alkyl groups; and R₅ and R₆ independently represent hydrogen, alkyl, phenyl, alkyl (optionally substituted) or R₅ and R₆ together with the nitrogen atom to which they are attached represent a saturated 3-7 membered heterocyclic ring (with a proviso); are disclosed which are antiinflammatory, antiallergic and immunomodulatory agents. Compositions containing these compounds and processes to prepare these compounds are also disclosed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

4-ALKYLAMINO-1-PHENOXYALKYL PIPERIDINES AS ANTIINFLAMMATORY, ANTIALLERGIC AND IMMUNOMODULATING AGENTS

This invention relates to novel substituted aminoalkyl-1-(aryloxyalkyl)piperidine compounds having therapeutic activity useful in treating conditions associated with inflammation, allergy or the immune system, to therapeutic compositions containing these novel compounds and to processes for preparing these novel compounds.

It is believed that, in response to an anti-inflammatory stimulus, phospholipase enzymes are activated leading to the release of arachidonic acid from phospholipids. Existing non-steroidal anti-inflammatory agents (NSAIA) are believed to act primarily by blocking the conversion of this released arachidonic acid into prostaglandins via the cyclooxygenase pathway of the arachidonic acid cascade. Many existing NSAIA are unsuitable for use by asthmatics. We have found a series of compounds which act to block the release of arachidonic acid from phospholipids. These compounds are indicated as useful antiinflammatory compounds with a potentially broader spectrum of activity than existing NSAIA, and potentially fewer gastro-intestinal side-effects. In addition the compounds may be useful in the treatment of asthma.

US 4,027,028 discloses that compounds of formula A




A

- 2 -

in which R_2 is a hydrogen or a lower alkyl radical; R_3 is the acyl residue of an organic alkylcarboxylic acid having up to 10 carbon atoms; R_4 is an optionally substituted phenyl radical; X is C_{2-3} alkylene radical
 5 optionally substituted by one or more lower alkyl radicals; n is 1 or 2; and Ar is inter alia optionally substituted phenyl; are useful for treating hypertension. Compounds of formula A in which R_3 represents hydrogen are disclosed as intermediates but
 10 no pharmacological activity is reported for these compounds.


US 4,348,401 discloses that compounds of formula B



in which R_1 is a hydrogen atom, an acyl radical or an optionally substituted aryl radical; R_2 is a
 15 hydroxymethyl radical, a cyano group, an amidino group which is optionally substituted by hydroxyl, a 1H-tetrazol-5-yl radical or a $-CO-R_3$ radical, R_3 is a hydroxyl group, a lower alkoxy radical or an amino group which is optionally substituted by a 1H-tetrazol-1-yl
 20 radical; X is an imino group or an oxymethyl radical; A is an alkylene radical containing 2 to 4 carbon atoms and B is a valency bond or a 4-hydroxypyrimidin-2,5-diyl radical; have an anti-allergic action, especially due to their strongly anti-histaminic action. The compounds
 25 are also alleged to have anti-oedematous and anti-phlogistic effectiveness.

GB 1,052,302 discloses that compounds of formula C

- 3 -




in which R_1 represents a straight- or branched-chain, saturated or unsaturated aliphatic hydrocarbon group containing at least two carbon atoms, a phenoxyalkyl group, an aralkyl group or an aralkenyl group; R_2 represents a straight- or branched-chain, saturated or unsaturated aliphatic hydrocarbon group, an aralkyl group, or an unsubstituted or substituted aromatic or aromatically unsaturated heterocyclic group; R_3 and R_4 which may be the same or different, each represents an alkyl or aralkyl group or, jointly, together with the adjacent nitrogen atom, represents a heterocyclic group which may contain a further hetero atom; and R_5 represents a hydrogen atom, or a lower alkyl group in the 3-position; exhibit analgesic activity and alleges that the compounds have spasmolytic, sedative and/or cough-soothing activity. However these compounds also exhibit cardiovascular activity which is undesirable.

EP 520,882 discloses the use of 4-aminomethyl-1-[2-(4-fluorophenoxy)ethyl]piperidine and 3-aminomethyl-1-[2-(2-methoxyphenoxy)ethyl]piperidine as chemical intermediates. No pharmacological activity is reported for these compounds.


EP 526,342 discloses the use of 4-(2-aminoethyl)-1-[2-(4-fluorophenoxy)ethyl]piperidine as a chemical intermediate. No pharmacological activity is reported for this compound.

Japanese Laid Open Patent Application No. 63-154683 (1988) discloses that certain pyridazinones are useful as heart stimulants. This application generically discloses intermediate compounds of formula D



- 5 in which R_2 , R_3 and R_4 independently represent hydrogen, a C_{1-5} alkoxy group or hydroxy; m and n are each whole numbers from 0-4 and X is oxygen, sulphur, $N(R_5)$ (in which R_5 is hydrogen or a C_{1-5} alkoxy group) or a single bond with the proviso that m is only 0 when X is a single bond. No specific examples are given of
 10 compounds of formula D in which X represents oxygen and no pharmacological activity is reported for compounds of formula D.

The present invention provides novel compounds of
 15 formula I



- and pharmaceutically acceptable salts thereof in which R_1 , R_2 and R_3 independently represent hydrogen, hydroxy, halo, a C_{1-6} alkyl group (optionally substituted by one or more halogen atoms) or a C_{1-6} alkoxy group
 20 (optionally substituted by one or more halogen atoms);

- 5 -

ALK¹ represents a C₂₋₆ alkylene chain optionally substituted by one or more C₁₋₂ alkyl groups;

Y represents a piperidine ring which is attached through nitrogen to ALK¹;

- 5 R₄ represents hydrogen or a C₁₋₄ alkyl group;

the broken line in --- represents a bond, so that a double bond connects Y and CR₄, or is absent and the free valency on Y is taken up by hydrogen and the free valency on CR₄ is taken up by hydrogen or a C₁₋₄ alkyl group;

10

ALK² is absent or represents a C₁₋₄ alkylene chain optionally substituted by one or more C₁₋₂ alkyl groups; and

- R₅ and R₆ independently represent hydrogen, a C₁₋₆ alkyl group, a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, halo or trifluoromethyl) or R₅ and R₆ together with the nitrogen atom to which they are attached represent a saturated 3-7 membered heterocyclic ring, optionally containing sulphur, oxygen or an additional nitrogen atom, wherein the ring is optionally substituted by one or more C₁₋₄ alkyl groups;
- 15
- 20


- excluding 4-aminomethyl-1-[2-(4-fluorophenoxy)ethyl]-piperidine, 4-(2-aminoethyl)-1-[2-(4-fluorophenoxy)-ethyl]piperidine and 3-aminomethyl-1-[2-(2-methoxyphenoxy)ethyl]piperidine.
- 25

It will be understood that a group containing a chain of three or more carbon atoms may be straight or

- 6 -

branched, for example propyl includes n-propyl and isopropyl and butyl includes n-butyl, sec-butyl, isobutyl and tert-butyl.

A preferred group of compounds of formula I is represented by formula II



and pharmaceutically acceptable salts thereof in which

R_1 , R_2 and R_3 independently represent hydrogen, hydroxy, halo (for example bromo, chloro or fluoro), trifluoromethyl, a C_{1-4} alkyl group (for example methyl, ethyl, propyl or butyl) or a C_{1-4} alkoxy group (for example methoxy, ethoxy, propoxy or butoxy);


$n = 2-4$ and $m = 1-3$;

R_5 represents hydrogen, a C_{1-4} alkyl group (for example methyl, ethyl, propyl or butyl) or a phenyl C_{1-4} alkyl group [(in which the phenyl ring is optionally substituted by one or more of the following: a C_{1-4} alkyl group, a C_{1-4} alkoxy group or halo) for example benzyl or phenethyl]; and

R_6 represents a C_{1-4} alkyl group (for example methyl, ethyl, propyl or butyl) or a phenyl C_{1-4} alkyl group [(in which the phenyl ring is optionally substituted by one or more of the following: a C_{1-4} alkyl group, a C_{1-4} alkoxy group or halo) for example benzyl or phenethyl];

or R₅ and R₆ together with the nitrogen atom to which they are attached represent a pyrrolidine ring, a piperidine ring or a morpholine ring.

A more preferred group of compounds of formula I is represented by formula III



and pharmaceutically acceptable salts thereof in which

R₁ represents hydrogen, halo, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group;

R₂ represents hydrogen, halo, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group;

R₃ represents hydrogen, a C₁₋₄ alkoxy group or hydroxy;

n = 2 and m = 1-3;

R₅ represents hydrogen, a C₁₋₄ alkyl group or a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group or halo); and

R₆ represents a C₁₋₄ alkyl group or a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group or halo);

or R₅ and R₆ together with the nitrogen atom to which they are attached represent a pyrrolidine ring, a piperidine ring or a morpholine ring.

The preferred values of R₁, R₂, R₃, R₅, R₆, and m in preferred compounds of formula III are listed below.

Preferably R₁ represents hydrogen, chloro, fluoro, methoxy or methyl. More preferably R₁ represents hydrogen, chloro, fluoro or methyl. Most preferably R₁ represents hydrogen or chloro.

10 Preferably R₂ represents hydrogen or chloro.

Preferably R₃ represents hydrogen, hydroxy or methoxy. More preferably R₃ represents hydrogen.

Preferably R₅ represents hydrogen, methyl, ethyl or propyl and R₆ represents methyl, ethyl, propyl or benzyl
15 or R₅ and R₆ together with the nitrogen atom to which they are attached represent a pyrrolidine ring or a piperidine ring. More preferably, R₅ represents methyl and R₆ represents methyl or benzyl or R₅ and R₆ together
20 with the nitrogen atom to which they are attached represent a piperidine ring. Most preferably, R₅ and R₆ both represent methyl.

Preferably m represents 1, 2 or 3. More preferably m represents 1 or 2. Most preferably m represents 1.

Specific compounds of formula I are:

- 25 4-aminomethyl-1-(2-phenoxyethyl)piperidine;
4-aminomethyl-1-[2-(4-chlorophenoxy)ethyl]piperidine;
4-aminomethyl-1-(3-phenoxypropyl)piperidine;

- 4-aminomethyl-1-[2-(4-methoxyphenoxy)ethyl]piperidine;
4-(dimethylaminomethyl)-1-[2-(4-methylphenoxy)ethyl]-
piperidine;
1-[2-(3-chlorophenoxy)ethyl]-4-(dimethylaminomethyl)-
5 piperidine;
4-(N-benzylaminomethyl)-1-(2-phenoxyethyl)piperidine;
4-(N-benzyl-N-methylaminomethyl)-1-(2-phenoxyethyl)-
piperidine;
4-(dimethylaminomethyl)-1-(2-phenoxyethyl)piperidine;
10 1-[2-(4-chlorophenoxy)ethyl]-4-(dimethylaminomethyl)-
piperidine;
4-(dimethylaminomethyl)-1-[2-(4-fluorophenoxy)ethyl]-
piperidine;
4-(dimethylaminomethyl)-1-[2-(4-methoxyphenoxy)ethyl]-
15 piperidine;
4-(dimethylaminomethyl)-1-(3-phenoxypropyl)piperidine;
4-(2-dimethylaminoethyl)-1-[2-(4-methylphenoxy)ethyl]-
piperidine;
4-(2-dimethylaminoethyl)-1-[2-(4-fluorophenoxy)ethyl]-
20 piperidine;
4-[2-(N-benzyl-N-methylamino)ethyl]-1-(2-phenoxyethyl)-
piperidine;
1-[2-(3-chlorophenoxy)ethyl]-4-(2-dimethylaminoethyl)-
piperidine;
25 1-(2-phenoxyethyl)-4-(2-piperidinoethyl)piperidine;
2-[2-[4-(2-dimethylaminoethyl)piperidino]ethoxy]phenol;
4-(2-dimethylaminoethyl)-1-(2-phenoxyethyl)piperidine;
1-(2-phenoxyethyl)-4-[2-(pyrrolidin-1-yl)ethyl]-
piperidine;
30 1-[2-(4-chlorophenoxy)ethyl]-4-(2-dimethylaminoethyl)-
piperidine;
4-(2-dimethylaminoethyl)-1-[2-(2-methoxyphenoxy)ethyl]-
piperidine;

- 4-(2-dimethylaminoethyl)-1-[2-(4-methoxyphenoxy)ethyl]-piperidine;
- 4-(3-dimethylaminopropyl)-1-(2-phenoxyethyl)piperidine;
- 4-(2-dimethylaminoethyl)-1-(3-phenoxypropyl)piperidine;
- 5 4-(2-dimethylaminoethyl)-1-(4-phenoxybutyl)piperidine;
- 4-[1-(4-chlorophenyl)ethylaminomethyl]-1-(2-phenoxyethyl)piperidine;
- 4-{2-[1-(4-chlorophenyl)ethylamino]ethyl}-1-(2-phenoxyethyl)piperidine;
- 10 4-(diethylaminomethyl)-1-(2-phenoxyethyl)piperidine;
- 4-(ethylaminomethyl)-1-(2-phenoxyethyl)piperidine;
- 4-(N-ethyl-N-methylaminomethyl)-1-(2-phenoxyethyl)-piperidine;
- 1-(2-phenoxyethyl)-4-(propylaminomethyl)piperidine;
- 15 4-(2-dimethylaminoethylidene)-1-(2-phenoxyethyl)-piperidine;
- 1-(2-phenoxyethyl)-4-[2-(pyrrolidin-1-yl)ethylidene]-piperidine;
- 4-[2-(N-benzyl-N-methylamino)ethylidene]-1-(2-phenoxyethyl)piperidine;
- 20 1-[2-(4-chlorophenoxy)ethyl]-4-(2-dimethylaminoethylidene)piperidine;
- 4-(2-dimethylaminoethylidene)-1-[2-(2-methoxyphenoxy)ethyl]piperidine;
- 25 4-(2-dimethylaminoethylidene)-1-[2-(4-methoxyphenoxy)ethyl]piperidine;
- 1-(2-phenoxyethyl)-3-(pyrrolidin-1-ylmethyl)piperidine;
- 1-(2-phenoxyethyl)-3-(piperidin-1-yl methyl)piperidine;
- (E)-3-(3-dimethylamino-1-propenyl)-1-(2-phenoxyethyl)-piperidine;
- 30 3-(3-dimethylaminopropyl)-1-(2-phenoxyethyl)piperidine;

and pharmaceutically acceptable salts thereof, in the form of individual enantiomers, racemates or other mixtures of enantiomers.

5 The compounds of formula I may form organic or inorganic salts, for example, the compounds of formula I may form acid addition salts with inorganic or organic acids, e.g. hydrochloric acid, hydrobromic acid, fumaric acid, tartaric acid, citric acid, sulphuric acid, 10 hydriodic acid, phosphoric acid, maleic acid, acetic acid, succinic acid, benzoic acid, pamoic acid, palmitic acid, dodecanoic acid and acidic amino acids such as glutamic acid. Some compounds of formula I may form base addition salts, for example, with alkali metal 15 hydroxides for example sodium hydroxide, or with aminoacids for example, lysine or arginine or with organic bases, for example meglumine. Certain compounds of formula I may exist in Zwitterionic form. It will be appreciated that all such salts, provided they are 20 pharmaceutically acceptable may be used in therapy in place of the corresponding compounds of formula I. Such salts are prepared by reacting the compound of formula I with a suitable acid or base in a conventional manner. Such salts may also exist in the form of solvates (for 25 example, hydrates).

It will be appreciated by those skilled in the art that certain compounds of formula I contain one or more chiral centres. Certain of the substituents R_1 , R_2 , R_3 , R_5 and R_6 may also contain at least one chiral centre, 30 for example when R_1 , R_2 , R_3 , R_5 and R_6 is sec-butyl.

When a compound of formula I, or a salt thereof, contains a single chiral centre it may exist in two enantiomeric forms. The present invention includes individual enantiomers and mixtures of those

enantiomers. The enantiomers may be obtained by methods known to those skilled in the art. Such methods typically include resolution via formation of diastereoisomeric salts or complexes which may be separated, for example, by crystallisation; resolution via formation of diastereoisomeric derivatives or complexes which may be separated, for example, by crystallisation, gas-liquid or liquid chromatography; selective reaction of one enantiomer by reaction with an enantiomer-specific reagent, for example, enzymatic esterification, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support such as silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation processes described above, a further step will subsequently be required to liberate the desired enantiomeric form. Alternatively, specific enantiomers may be synthesised by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation.

When a compound of formula I, or a salt thereof, contains more than one chiral centre it may exist in diastereoisomeric forms. The diastereoisomeric pairs may be separated by methods known to those skilled in the art, for example, chromatography or crystallisation and the individual enantiomers within each pair may be separated as described above. The present invention includes each diastereoisomer of compounds of formula I and mixtures thereof.

- 13 -

Certain compounds of formula I may exist in more than one geometric isomeric form for example when a double bond is present between Y and C(R⁴). The present invention includes each geometric isomer and mixtures thereof.

Certain compounds of formula I and their salts may exist in more than one crystal form and the present invention includes each crystal form and mixtures thereof. Certain compounds of formula I may also exist in the form of solvates, for example hydrates, and the present invention includes each solvate and mixtures thereof.

The present invention also provides pharmaceutical compositions comprising a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof (including the three excluded compounds) together with a pharmaceutically acceptable diluent or carrier. Such pharmaceutical compositions may be used in the treatment of inflammatory and/or allergic diseases.

As used hereinafter, the term "active compound" denotes a compound of formula I or a pharmaceutically acceptable salt thereof. In therapeutic use the active compound may be administered orally, rectally, parenterally, topically, ocularly, aurally, nasally, intravaginally or to the buccal cavity, to give a local and/or a systemic effect. The active compounds may be administered in a prophylactic manner. Thus the therapeutic compositions of the present invention may take the form of any of the known pharmaceutical compositions for such methods of administration. The compositions may be formulated in a manner known to those skilled in the art so as to give a controlled

- 14 -

- release, for example rapid release or sustained release, of the compounds of the present invention. Pharmaceutically acceptable carriers suitable for use in such compositions are well known in the art of pharmacy.
- 5 The compositions of the invention suitably contain 0.1-90% by weight of active compound. The compositions of the invention are generally prepared in unit dosage form.

- Compositions for oral administration are the
- 10 preferred compositions of the invention and these are the known pharmaceutical forms for such administration, for example tablets, capsules, granules, syrups and aqueous or oily suspensions. The excipients used in the preparation of these compositions are the excipients
- 15 known in the pharmacists' art.

- Tablets may be prepared from a mixture of the active compound with fillers, for example, lactose or calcium phosphate, disintegrating agents, for example maize starch, lubricating agents, for example magnesium
- 20 stearate, binders for example microcrystalline cellulose or polyvinyl pyrrolidone and other optional ingredients known in the art to permit tableting the mixture by known methods. The tablets may, if desired, be coated using known methods and excipients which may include
- 25 enteric coating using for example hydroxypropylmethyl-cellulose phthalate.

- The tablets may be formulated in a manner known to those skilled in the art so as to give a sustained release of the compounds of the present invention. Such
- 30 tablets may if desired be provided with enteric coatings by known methods, for example by the use of cellulose acetate phthalate.

- 15 -

Similarly capsules, for example hard or soft gelatin capsules containing the active compound with or without added excipients, may be prepared by known methods and, if desired, provided with enteric coatings in a known manner. The contents of the capsule may be formulated using known methods to give sustained release of the active compound. Enteric coated compositions of the invention may be advantageous, depending on the nature of the active compound. The tablets and capsules may conveniently each contain 1-1000 mg (for example 10 mg, 50 mg, 100 mg, 200 mg, 400 mg, 600 mg or 800 mg) of the active compound. Other compositions for oral administration include, for example, aqueous suspensions containing the compound of formula I in an aqueous medium in the presence of a non-toxic suspending agent such as sodium carboxymethylcellulose, and oily suspensions containing a compound of the present invention in a suitable vegetable oil, for example sunflower oil.

The active compound may be formulated into granules with or without additional excipients. The granules may be ingested directly by the patient or they may be added to a suitable liquid carrier (for example water) before ingestion. The granules may contain disintegrants (for example a pharmaceutically acceptable effervescent couple formed from an acid and a carbonate or bicarbonate salt) to facilitate dispersion in the liquid medium.

Compositions of the invention suitable for rectal administration are the known pharmaceutical forms for such administration, for example suppositories with hard fat, semi-synthetic glycerides or polyethylene glycol bases.

- 16 -

Compositions of the invention suitable for parenteral administration are the known pharmaceutical forms for such administration, for example sterile suspensions in aqueous and oily media or sterile solutions in a suitable solvent.

Compositions for topical administration may comprise a matrix in which the active compound is dispersed so that it is held in contact with the skin in order to administer the active compound transdermally. Alternatively the active compound may be dispersed in a cream, gel or ointment base or applied in the form of a spray.

Compositions of the invention suitable for inhalation via the mouth and/or the nose are the known pharmaceutical forms for such administration, for example aerosols, nebulised solutions or powders. Metered dose systems, known to those skilled in the art, may be used.

Compositions suitable for application to the buccal cavity include slow dissolving tablets, troches, chewing gum, gels, pastes, powders, mouthwashes or rinses.

The compounds of the present invention may also be administered by continuous infusion either from an external source, for example by intravenous infusion or from a source of the compound placed within the body. Internal sources include implanted reservoirs containing the compound to be infused which is continuously released for example by osmosis and implants which may be a) liquid such as an oily solution or suspension of the compound to be infused for example in the form of a very sparingly water-soluble derivative such as a dodecanoate salt or b) solid in the form of an implanted

- 17 -

support for example a synthetic resin or waxy material for the compound to be infused. The support may be a single body containing all the compound or a series of several bodies each containing part of the compound to be delivered.

In some formulations it may be beneficial to use the compounds of the present invention in the form of particles of very small size, for example as obtained by fluid energy milling.

10 In the compositions of the present invention the active compound may, if desired, be associated with other compatible pharmacologically active ingredients, for example:

15 a) an analgesic (e.g. in treatment of rheumatoid arthritis), b) a β_2 agonist (e.g. in treatment of asthma) and c) a non-sedating antihistamine (e.g. in treatment of other allergic conditions).

The pharmaceutical compositions containing a therapeutically effective amount of a compound of formula I may be used to treat inflammatory and/or allergic conditions in human beings. In such treatment the amount of the compound of formula I administered per day is in the range 0.1 to 3000 mg. Specific compounds which may be incorporated into the compositions of this invention are the novel compounds disclosed above.

The therapeutic activity of compounds of formula I has been demonstrated by means of tests on standard laboratory animals. Such tests include, for example, the oral administration of the compounds to rats in which an inflammatory condition is induced. Thus, compounds of formula I are useful for the treatment of inflammatory conditions in mammals. Whilst the precise

amount of active compound administered will depend on a number of factors, for example the age of the patient, the severity of the condition and the past medical history and always lies within the sound discretion of the administering physician, a suitable dose for enteral administration to mammals, including humans, is generally within the range 0.01-80 mg/kg/day, more usually 0.2-40 mg/kg/day given in single or divided doses. For parenteral administration, a suitable dose is generally within the range 0.01-80 mg/kg/day, more usually 0.2-40 mg/kg/day given in single or divided doses or by continuous infusion. Oral administration is preferred.

Compounds of formula I and pharmaceutically acceptable salts thereof (including the three excluded compounds) are indicated for use as medicaments particularly in the treatment of inflammatory and/or allergic conditions for example musculoskeletal disorders for example: rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, muscle trauma, gout, ankylosing spondylitis, tendonitis and bursitis; respiratory disorders for example: asthma and rhinitis; gastrointestinal disorders for example: gastritis, Crohn's disease, ulcerative colitis and other inflammatory diseases of the bowel; diseases of the oral cavity for example: periodontitis and gingivitis; cutaneous disorders for example: psoriasis, urticaria, allergic skin diseases, burns, ocular inflammation and iritis; or Alzheimer's disease. Compounds of formula I and salts thereof may also be useful as analgesics and/or anti-pyretic agents.

Accordingly, in another aspect, the present invention also includes a method of treating inflammatory and/or allergic conditions comprising the

administration of a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof (including the three excluded compounds) to a mammal in need thereof.

- 5 While the precise mechanism of action of the compounds of formula I is unknown at present, it is believed that the pharmacological effects arise from the ability of these compounds to inhibit the release of arachidonic acid from phospholipids. Consequently, in a
10 preferred aspect, the present invention provides a method of treating inflammatory and/or allergic conditions comprising the administration of a therapeutically effective amount of an arachidonic acid release inhibitor of formula I or a pharmaceutically
15 acceptable salt thereof (including the three excluded compounds) to a mammal in need thereof.

- In yet another aspect, the present invention provides the use of a compound of formula I or a pharmaceutically acceptable salt thereof (including the
20 three excluded compounds) in the manufacture of a medicament for use in the treatment of an inflammatory and/or allergic condition.

- The compounds of formula I may also be indicated for use as immunomodulatory agents, generally as
25 immunosuppressants, but some compounds, in certain disease states, may exhibit immunostimulant activity. The compounds according to the invention may be useful in the treatment of diseases resulting from an aberrant immune reaction. Thus the pharmaceutical compositions
30 containing a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof (including the three excluded compounds) may be used to treat diseases with an immunological

- 20 -

association for example tissue rejection, such as kidney rejection; autoimmune diseases, such as thyroiditis and type 1 diabetes; cutaneous disorders, such as contact sensitivity, eczema and psoriasis; neoplasia, such as melanoma; and HIV infection.

In such treatment the amount of the compound of formula I administered per day will be such as to give a therapeutic effect and is generally in the range 0.1 to 2000 mg, preferably 1 to 500 mg.

Accordingly, in another aspect, the present invention also includes a method of treating diseases with an immunological association, comprising the administration of a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof (including the three excluded compounds) to a mammal in need thereof.

In yet another aspect, the present invention provides the use of a compound of formula I or a pharmaceutically acceptable salt thereof (including the three excluded compounds) in the manufacture of a medicament for use in the treatment of diseases with an immunological association.

The therapeutic activity of compounds falling within formula I may be demonstrated by means of in vitro and in vivo tests. Such tests include, for example, the in vitro mixed lymphocyte reaction and an in vivo rat Graft versus Host (GVH) test. Thus, compounds of formula I may be useful as immunomodulatory agents.


The compounds of formula I and pharmaceutically acceptable salts thereof may also be indicated for use

- 21 -


as disease-modifying antirheumatic agents. Such activity may be demonstrated by means of in vivo tests such as the antigen induced mouse arthritis test as described in WO 93/13097.

5 Processes for the preparation of compounds of formula I will now be described. These processes form a further aspect of the present invention. Unless otherwise stated the processes are carried out at atmospheric pressure.

10 Compounds of formula I may be prepared by reacting a compound of formula IV




in which R₁, R₂, R₃ and ALK¹ are as previously defined and L₁ represents a leaving group, for example halo or tosyl, with a compound of formula V




15 in which Y, R₄, ALK², R₅ and R₆ are as previously defined, optionally in the presence of an inert organic liquid which is preferably a solvent for the reactants, e.g. N,N-dimethylformamide, at a temperature in the range from 0°C up to the boiling point of the inert organic liquid, e.g. 0-250°C, preferably by heating at a temperature in the range 20-150°C, optionally in the presence of a base e.g. triethylamine.

Compounds of formula I in which R_5 represents a group other than hydrogen may be prepared by reacting a compound of formula VI



in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 and ALK^2 are as initially defined and $R_{12}CO$ represents a group which on reduction yields a group R_5 of formula $-CH_2R_{12}$ and R_{11} represents R_6 (or $R_{12}CO$ to give compounds in which R_5 and R_6 are identical), with a reducing agent, for example lithium aluminium hydride, preferably in the presence of an inert organic liquid which is preferably a solvent for the compound of formula VI, for example tetrahydrofuran, at a temperature in the range from -50°C up to the boiling point of the inert organic liquid e.g. -50 to 200°C , preferably at a temperature in the range -10 to 150°C .

Compounds of formula I in which R_5 represents hydrogen may be prepared by reacting a compound of formula VII




in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 and ALK^2 are as initially defined and $-N=R_{13}$ represents a group which on reduction yields $-NHR_6$, for example benzylimino, with a

- 23 -


reducing agent, for example sodium borohydride, preferably in the presence of an organic liquid which is a solvent for the compound of formula VII, for example an alcohol e.g. ethanol, at a temperature in the range
 5 0-200°C, preferably by heating at a temperature in the range 20-150°C. "

Compounds of formula I in which R_5 represents hydrogen may be prepared by deprotecting a compound of formula VIII




10 in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 , R_6 and ALK^2 are as initially defined and PG_1 represents an amine protecting group for example benzoyl e.g. by hydrolysis. Suitable methods for protecting and deprotecting amines may be found in the textbook "Protective Groups in Organic
 15 Synthesis" by T.W. Greene, John Wiley and Sons, 1981.

Compounds of formula I in which (CR_4) is connected to Y by a double bond may be prepared by reacting a compound of formula IX




20 in which R_1 , R_2 , R_3 , ALK^1 and Y are as initially defined, with a Wittig reagent, for example (a) a compound of formula Xa

- 24 -




in which ALK^2 , R_4 , R_5 and R_6 are as initially defined and X represents halo,

or (b) a compound of formula Xb



- in which ALK^2 , R_4 , R_5 and R_6 are as initially defined and R_{16} represents an alkyl group, in which Xa or Xb has been pretreated with a base, e.g. n-butyllithium, in an inert organic liquid e.g. tetrahydrofuran, and the mixture combined in an inert organic liquid e.g. tetrahydrofuran, at a temperature in the range from 0°C up to the boiling point of the inert organic liquid, e.g. 0-200°C, preferably by heating at a temperature in the range 20-150°C.


Compounds of formula I may be prepared by reducing compounds of formula XI



- 25 -


in which R_1 , R_2 , R_3 , Y , R_4 , ALK^2 , R_5 and R_6 are as initially defined and $-ALK^3-CO-$ represents a group which on reduction yields a group of formula $-ALK^1-$, for example using borane, in an inert organic liquid which is preferably a solvent for XI, for example tetrahydrofuran, at a temperature in the range from $0^\circ C$ up to the boiling point of the inert organic liquid e.g. $0-200^\circ C$, preferably by heating at a temperature in the range $20-150^\circ C$.

Compounds of formula I in which R_5 and R_6 represent hydrogen may be prepared by hydrolysing compounds of formula XII



in which R_1 , R_2 , R_3 , ALK^1 , R_4 and ALK^2 are as initially defined and PG_2 represents an amine protecting group, for example dimethylaminomethylene or 4-chloro- α -methylbenzylidene, preferably in the presence of a base, e.g. aqueous ammonia, or an acid, e.g. a mineral or an organic acid, e.g. hydrochloric acid or acetic acid, at a temperature in the range $0-250^\circ C$, preferably by heating at a temperature in the range of $20-100^\circ C$.

Compounds of formula I may be prepared by reducing a compound of formula XIII




in which R_1 , R_2 , R_3 , Y , R_4 , R_5 and R_6 are as initially defined and $-(ALK^3)-CO-$ and $(ALK^4)-CO-$ represent groups which on reduction give ALK^1 and ALK^2 respectively, for example using borane, in an inert organic liquid which

5 is preferably a solvent for the compound of formula XIII, for example tetrahydrofuran, at a temperature in the range from $0^\circ C$ up to the boiling point of the inert organic liquid, e.g. $0-200^\circ C$, preferably by heating at a temperature in the range $20-150^\circ C$. It will be

10 appreciated by those skilled in the art that if an olefinic double bond is present in XIII that this bond may also be reduced.

Compounds of formula I in which $---CR_4-$ represents methylene, ALK^2 is absent and R_5 and R_6 represent

15 hydrogen may be prepared by reacting a compound of formula XIV




in which R_1 , R_2 , R_3 , ALK^1 are as initially defined, ALK^6 is absent and Y^1 represents a piperidine ring, or a pyridinium ring having a halide counter ion, with a

20 reducing agent, for example lithium aluminium hydride or hydrogen in the presence of a catalyst, e.g. palladium on charcoal, in an inert organic liquid which is

- 27 -


preferably a solvent for the compound of formula XIV,
for example tetrahydrofuran, at a temperature in the
range from 0°C up to the boiling point of the inert
organic liquid, e.g. 0-200°C, preferably by heating at a
5 temperature in the range 20-150°C.

Compounds of formula I in which the broken line is
absent, the free valency on Y is taken up by hydrogen,
the free valency on CR₄ is taken up by hydrogen or a
C₁₋₄ alkyl group, R₅ and R₆ each represent hydrogen and
10 ALK² represents a C₁₋₄ alkylene chain may be prepared by
reacting a compound of formula XIV




in which -(ALK⁶)-CN represents a group which on
reduction gives -CH(R₄)-(ALK²)-NH₂ and Y¹ represents a
piperidine ring, or a pyridinium ring having a halide
counter ion, with a reducing agent, for example lithium
15 aluminium hydride or hydrogen in the presence of a
catalyst, e.g. palladium on charcoal, in an inert
organic liquid which is preferably a solvent for the
compound of formula XIV, for example tetrahydrofuran, at
20 a temperature in the range from 0°C up to the boiling
point of the inert organic liquid, e.g. 0-200°C,
preferably by heating at a temperature in the range
20-150°C.

Compounds of formula I may be prepared by reacting
25 a compound of formula XV




in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 and ALK^2 are as initially defined and L_6 represents a leaving group, for example halo, with a compound of formula XXX




in which R_5 and R_6 are as initially defined, optionally
 5 in the presence of an inert organic liquid which is preferably a solvent for the reactants, for example N,N-dimethylformamide or tetrahydrofuran, at a temperature in the range from 0°C up to the boiling point of the inert organic liquid e.g. $0\text{-}250^\circ\text{C}$, preferably by heating
 10 at a temperature in the range $20\text{-}150^\circ\text{C}$.

Compounds of formula I may be prepared by reacting a compound of formula XVI




in which R_1 , R_2 and R_3 are as initially defined with a compound of formula XVIII



in which ALK^1 , Y , R_4 , ALK^2 , R_5 and R_6 are as initially defined, in the presence of a dialkyl azodicarboxylate, for example diethyl azodicarboxylate, and a phosphorus (III) reagent for example, triphenylphosphine, optionally in the presence of an inert organic liquid which is preferably a solvent, for example tetrahydrofuran, at a temperature in the range from 0°C up to the boiling point of the inert organic liquid, for example $0-250^\circ\text{C}$, preferably by heating at a temperature in the range $0-150^\circ\text{C}$.


Compounds of formula I in which R_5 and R_6 each represent hydrogen and ALK^2 is absent may be prepared by reacting compounds of formula XL




in which R_1 , R_2 , R_3 , ALK^1 , Y and R_4 are as initially defined with a reducing agent, for example iron and ammonium chloride solution or hydrogen in the presence of a catalyst. It will be appreciated by those skilled in the art that if an olefinic double bond is present in XL that this bond may also be reduced or left intact, as desired, by the choice of reducing agent.

Compounds of formula I may be prepared by reacting a compound of formula V

- 30 -




in which Y, R₄, ALK², R₅ and R₆ are as initially defined with a compound of formula XLI




in which R₁, R₂ and R₃ are as initially defined and -(ALK⁷)-CHO represents a group which on reaction and
 5 reduction yields a group of formula ALK¹, optionally in the presence of an inert organic liquid which is preferably a solvent for V and XLI, for example toluene, at a temperature in the range from 0°C up to the boiling point of the inert organic liquid e.g. 0-250°C,
 10 preferably by heating at a temperature in the range 20-150°C, followed by reaction of the product with a reducing agent, for example sodium borohydride or sodium cyanoborohydride, optionally in the presence of an inert organic liquid which is preferably a solvent for the
 15 reactants, e.g. ethanol, at a temperature in the range 0-250°C, preferably by heating at a temperature in the range 20-150°C.

Compounds of formula I may be prepared by reacting a compound of formula XVI with a compound of formula
 20 XLII




- in which ALK^1 , Y , R_4 , ALK^2 , R_5 and R_6 are as initially defined and L_8 represents a leaving group, for example halo or tosyl, optionally in the presence of an inert organic liquid, which is preferably a solvent for the reactants, for example N,N-dimethylformamide or an alcohol, at a temperature in the range 0-250°C, preferably by heating at a temperature in the range 20-150°C, optionally in the presence of a base, for example sodium hydroxide.

- 10 Compounds of formula IV may be prepared by methods known to those skilled in the art for example by reacting a compound of formula XVI with a compound of formula XVII



- 15 in which ALK^1 is as initially defined and L_1 and L_2 independently represent a leaving group which may be the same or different provided that L_2 is more labile than L_1 , for example halo or tosyl.


Compounds of formula V may be prepared by hydrolysing compounds of formula XIX



- 32 -


- in which Y, R₄, ALK², R₅ and R₆ are as initially defined and R₁₄CO represents a hydrolysable acyl group, for example acetyl or benzoyl, for example by reaction with dilute hydrochloric acid at a temperature in the range 0-250°C, preferably by heating at a temperature in the range 20-150°C, optionally followed by basification.

Compounds of formula VI may be prepared by reacting a compound of formula XX




- in which Y, R₄, ALK², R₁₁ and R₁₂CO are as initially defined, with a compound of formula IV in an analogous manner to the preparation of a compound of formula I from a compound of formula IV and a compound of formula V.

- Compounds of formula VI may also be prepared by reacting a compound of formula I in which R₅ and/or R₆ represents hydrogen with a compound of formula XXI




in which R₁₂CO is as initially defined and L₃ represents a leaving group, for example halo, by methods known to those skilled in the art.

- Compounds of formula VII may be prepared by reacting a compound of formula XXII



in which Y, R₄, ALK² and -N=R₁₃ are as initially defined, with a compound of formula IV in an analogous manner to the preparation of a compound of formula I from a compound of formula IV and a compound of formula V.

Compounds of formula VIII may be prepared in a similar manner to compounds of formula VII by reacting a compound of formula XXIII



in which Y, R₄, ALK², PG₁ and R₆ are as initially defined with a compound of formula IV.


Compounds of formula IX may be prepared by reacting a compound of formula XXIV



in which Y is as initially defined, with a compound of formula IV optionally in the presence of an inert organic liquid which is preferably a solvent for the reactants, at a temperature in the range 0-250°C optionally in the presence of a base, for example triethylamine or sodium bicarbonate.


Compounds of formula Xa and Xb may be prepared by methods known to those skilled in the art for example Xa

may be prepared by reacting triphenylphosphine with a compound of formula XXV




in which R_4 , ALK^2 , R_5 and R_6 are as initially defined.

Compounds of formula XI may be prepared by reacting
5 a compound of formula XXVI




in which R_1 , R_2 , R_3 and ALK^3 are as initially defined
and L represents a leaving group, for example halo, with
a compound of formula V for example in the presence of a
base, e.g. triethylamine, in an inert organic liquid,
10 e.g. dichloromethane at a temperature in the range from
0°C up to the boiling point of the inert organic liquid,
e.g. 0-200°C, preferably by heating at a temperature in
the range 20-150°C.

Compounds of formula XI may also be prepared by
15 reacting a compound of formula XXVII




- 35 -

in which R_1 , R_2 , R_3 , ALK^3 , Y , R_4 and ALK^2 are as initially defined and Hal represents chloro, bromo or iodo with a compound of formula XXX




in which R_5 and R_6 are as initially defined, for example
 5 in an inert organic liquid, e.g. tetrahydrofuran, at a temperature in the range 0-150°C in the presence of a base, for example triethylamine, preferably by heating at a temperature in the range 20-150°C. It will be appreciated by those skilled in the art that this
 10 reaction may give rise to quaternary salt by-products from which the desired product may be separated by known techniques, e.g. distillation.

Compounds of formula XII may be prepared by reacting a compound of formula XXXIX




in which R_4 , ALK^2 and PG_2 are as initially defined with a compound of IV for example in the presence of a base, e.g. triethylamine in an inert organic liquid, e.g. dichloromethane or toluene, at a temperature in the
 20 range from 0°C up to the boiling point of the inert organic liquid, e.g. 0-200°C, preferably by heating at a temperature in the range 20-150°C.

Compounds of formula XIII may be prepared by reacting a compound of formula XXIX



in which R_1 , R_2 , R_3 , ALK^3 , Y , R_4 , ALK^4 and Hal are as initially defined with a compound of formula XXX by methods known to those skilled in the art.

- Compounds of formula XIV in which Y^1 represents
 5 piperidine may be prepared by reacting a compound of formula XXVIII




- in which R_1 , R_2 , R_3 , ALK^1 , Y and ALK^6 are as initially defined and L_7 represents a leaving group, for example halo or tosyl, with a cyanide salt for example potassium
 10 cyanide, by methods known to those skilled in the art.

- Compounds of formula XIV in which Y^1 represents a piperidine ring may also be prepared by reducing compounds of formula XIV, in which Y^1 represents a pyridinium ring with a halide counter ion, by methods
 15 known to those skilled in the art for example by reacting with sodium borohydride in a solvent e.g. ethanol.

- Compounds of formula XIV in which Y^1 represents a pyridinium ring having a halide counter ion may be
 20 prepared by alkylation of an appropriate cyanopyridine or cyanoalkylpyridine, for example with a compound of

formula IV, by methods known to those skilled in the art.

Compounds of formula XV may be prepared from compounds of formula XXXVIII




5 in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 and ALK^2 are as initially defined by methods known to those skilled in the art.

Compounds of formulae XVI, XVII and XXI are commercially available or may be prepared by methods
10 known to those skilled in the art.


Compounds of formula XVIII may be prepared from compounds of formula V by methods known to those skilled in the art for example by hydroxyalkylation.

Compounds of formula XIX may be prepared by
15 reacting a compound of formula XXXI



in which R_{14} , Y , R_4 and ALK^2 are as initially defined and L_5 represents a leaving group, for example halo, with a compound of formula XXX in an analogous manner to the preparation of a compound of formula XI from a
20 compound of formula XXVII and a compound of formula XXX.


Compounds of formula XX may be prepared by acylating a compound of formula XXXII



in which Y, R₄, ALK² and R₆ are as initially defined, or by diacylating a compound of formula XXXII in which R₆ represents hydrogen, by methods known to those skilled in the art.

Compounds of formula XXII may be prepared from compounds of formula XXXII by methods known to those skilled in the art, for example by reacting a compound of formula XXXII, in which R₆ is hydrogen, with an aldehyde of formula HR₁₃=O.


Compounds of formula XXIII may be prepared by protecting compounds of formula XXXII



in which Y, R₄, ALK² and R₆ are as initially defined by methods known to those skilled in the art.

Compounds of formula XXIV, XXV, XXVI, XXX and XLI are commercially available or may be prepared by methods known to those skilled in the art.


Compounds of formula XXVII may be prepared by reacting a compound of formula XXXIII



in which R_1 , R_2 , R_3 , ALK^3 , Y , R_4 and ALK^2 are as initially defined, with a halogenating agent, e.g. thionyl chloride, at a temperature in the range 0-200°C in the presence or absence of an inert organic liquid, e.g. chloroform, preferably by heating at a temperature in the range 20-150°C.


Compounds of formula XXVIII may be prepared in an analogous manner to compounds of formula XV.

Compounds of formula XXIX may be prepared by
10 halogenating a compound of formula XXXIV



in which R_1 , R_2 , R_3 , ALK^3 , Y , R_4 and ALK^4 are as initially defined and R_{15} represents hydrogen for example by reaction with thionyl chloride at a temperature in the range 0-200°C in the presence or the
15 absence of an inert organic liquid, e.g. chloroform, preferably by heating at a temperature in the range of 20-150°C.


Compounds of formula XXXI may be prepared from compounds of formula XXXV



in which R_{14} , Y , R_4 and ALK^2 are as initially defined, for example, by reaction with the halogenating agent, for example thionyl chloride at a temperature in the range 0-200°C in the presence or the absence of an inert organic liquid, e.g. chloroform, preferably by heating at a temperature in the range of 20-100°C.


Compounds of formula XXXII may be prepared by methods analogous to those described for the preparation of compounds of formula V.

Compounds of formula XXXIII may be prepared by reacting a compound of formula XXXVI



in which Y , R_4 and ALK^2 are as initially defined, with a compound of formula XXVI for example, by reaction in the presence of a base, e.g. triethylamine, in an inert organic liquid, e.g. dichloromethane, at a temperature in the range from 0°C up to the boiling point of the inert organic liquid, e.g. 0-200°C, preferably by heating in the range 20-150°C.

Compounds of formula XXXIV may be prepared by reacting a compound of formula XXXVII



in which Y, R₄ and ALK⁴ are as initially defined and R₁₅ represents a C₁₋₄ alkyl group with a compound of formula XXVI by reaction in the presence of a base, e.g. triethylamine, in an inert organic liquid, e.g. methylene chloride, at a temperature in the range of 0°C up to the boiling point of the inert organic liquid, e.g. 0-200°C, preferably by heating to a temperature in the range 20-150°C to form a compound of formula XXXIV in which R₁₅ represents a C₁₋₄ alkyl group and then hydrolysing by reaction with sodium hydroxide in an aqueous non-alcoholic solvent to give a compound of formula XXXIV in which R₁₅ represents hydrogen.

Compounds of formula XXXV may be prepared by acylating compounds of formula XXXVI by methods known to those skilled in the art for example by reacting with an acylating agent optionally in the presence of a base, e.g. triethylamine, and then reacting with base at a temperature in the range 0-100°C to cleave any O-acylated material.

Compounds of formula XXXVI and XXXVII are commercially available or may be prepared by methods known to those skilled in the art.

Compounds of formula XXXVIII may be prepared by reaction of a compound of formula XXXVI with a compound of formula IV by methods known to those skilled in the art.

Compounds of formula XXXIX may be prepared by protecting compounds of formula HY---CR₄-(ALK²)-NH₂ (which may be prepared by methods analogous to those described for the preparation of compounds of formula V) by methods known to those skilled in the art.

Compounds of formula XL may be prepared by methods known to those skilled in the art for example by reacting a compound of formula IX with a compound of formula $H_2C(R_4)-NO_2$, for example using the Knoevenagel condensation.

Compounds of formula XLII may be prepared by methods known to those skilled in the art, for example from compounds of formula XVIII.

It will be appreciated by those skilled in the art that the substituents specified for R_1 , R_2 and R_3 may be interconverted by methods known to those skilled in the art. Also certain compounds of formula I may be converted into other compounds of formula I as indicated below.

Compounds of formula I in which R_5 and/or R_6 represents a group other than hydrogen may be prepared by alkylating a compound of formula I in which R_6 and/or R_5 , respectively, represents hydrogen either

a) directly using an alkylating agent of formula R_5L_3 or R_6L_3 in which L_3 represents a leaving group for example halo in the presence of an organic liquid which is preferably a solvent for the reactants at a temperature in the range 0-150°C optionally in the presence of a base or

b) by reductive alkylation, for example using formaldehyde and formic acid, at a temperature in the range 0-200°C, preferably by heating at a temperature in the range 20-150°C, optionally in the presence of an inert organic liquid, which is preferably a solvent for the reactants.

- 43 -

Compounds of formula I in which --- represents a single bond may be prepared by hydrogenating a compound of formula I in which --- represents a double bond, for example by reaction with hydrogen in the presence of a catalyst e.g. palladium on charcoal, in the presence of an inert organic liquid, which is preferably a solvent for the starting compound of formula I, for example an alcohol at a temperature in the range 0-250°C preferably 15-100°C, preferably at a pressure of 1 atmosphere.

10 It will be appreciated by those skilled in the art that alkylation reactions may give rise to quaternary salt by-products from which the desired product may be separated by known techniques for example distillation.

15 It will be appreciated by those skilled in the art that in cases where a substituent is identical with, or similar to, a functional group which is being modified in one of the above processes that these substituents will require protection before the process is undertaken, followed by deprotection after the process.

20 Otherwise competing side-reactions will occur. Alternatively, another of the processes described above, in which the substituent does not interfere, may be used.

Compounds of formula II and III may be prepared by processes analogous to those described for the preparation of compounds of formula I.

The compounds of formula I are antiinflammatory agents and may show therapeutic activity at a dose of 200 mg/kg or lower in standard laboratory animals. The therapeutic activity of compounds of formula I has been demonstrated by Test A and one or more of Tests B, C and D.

Test A was carried out in the following way:-

Inhibition of Arachidonic Acid Release from Zymosan
Stimulated Macrophages

Female MF1 mice (weighing 20 to 25 g) were killed
5 using a rising concentration of CO₂. The mice were laid
on their backs and the abdomens wiped with 70% alcohol.
The skin was pulled back, exposing the peritoneal wall.
Medium A (5 ml) (see below) was injected into the
peritoneal cavity of each mouse followed by
10 approximately 1 ml of air using a 20 ml syringe and a
21G x 40 mm needle in order to form a suspension of
macrophage cells. The medium and cells were then
removed using a 19G x 40 mm needle. The resulting
suspension was returned to a sterile beaker kept on ice.
15 The extracts from all the mice were pooled and this
pooled cell suspension was counted using a Coulter
counter and adjusted to a final cell count of $1 - 1.3 \times 10^6$ cells/ml prior to labelling with [³H]-
arachidonic acid. Typically five mice provided
20 sufficient cells for each multiwell plate.

Sufficient [³H]-arachidonic acid in ethanol to give
a final concentration of 1.6 µCi/ml (equivalent to
40 µCi/plate) was blown to dryness under nitrogen. The
arachidonic acid was then resuspended in 1 or 2 ml of
25 the cell suspension which was then mixed with the
remainder of the cell suspension in a centrifuge bottle.
The labelled cell suspension was then plated out into
sterile plastic 96 flat-bottomed well plates (250 µl per
well) and incubated overnight at 37°C in a moist
30 atmosphere of 5% CO₂, 95% air.

The following day, non-adherent cells were removed
by washing 3 times with sterile phosphate buffered

saline (PBS). The adherent peritoneal macrophages were then cultured for a further 24 hours in the presence or absence of drugs, in medium B (see below) at 37° in a 5% CO₂ atmosphere in order to measure the effects of drugs on the spontaneous release of arachidonic acid in the absence of stimulus. After this incubation, supernatants were removed to give medium 1 and stored in sealed multi-well plates at 4°C prior to scintillation counting. Drugs which potentiated spontaneous release of arachidonic acid (125% of controls) were deemed to be toxic at the concentration at which this phenomenon occurred. The supernatants were replaced by fresh medium C containing fresh drug and a stimulus. Three drugs were tested at six concentrations (100, 50, 20, 10, 5 and 1 µM) in replicates of four on each plate. The other wells contained controls consisting of a positive control (e.g. dexamethasone), medium (B) only and medium C only.

Incubation was then continued for a further 5 hours, whereupon the supernatants were collected to give medium 2 and the adherent cells washed with PBS. The cells were then lysed with 100 µl of 0.1% TRITON® X100 in a 0.1% solution of bovine serum albumin in 0.9% saline and mechanically disrupted to give cell lysates. These supernatants (medium 2) and cell lysates (Cells) were also stored in sealed multi-well plates at 4°C prior to scintillation counting. 200 µl aliquots of media, or 100 µl aliquots of cells were counted using 2 ml of OPTIPHASE "HIGH SAFE" (Trademark of LKB) as scintillant.

Calculation of results

The percentage of arachidonic acid released was calculated using the mean values for each group of 4 wells in the following equation.

$$\% \text{ Release} = \frac{\text{cpm in medium 2}}{\text{cpm in medium 2} + \text{cpm in cell lysate}} \times 100$$

cpm = counts per minute

The value for the arachidonic acid release in the absence of stimulus (spontaneous, cpm of medium 2) from cells which had been exposed to neither stimulus nor drug was subtracted from all equivalent values (cpm media 2, stimulated with or without drug) to give the net stimulated release. The percentage inhibition of arachidonic acid release caused by a drug may then be calculated using the following equation.

$$\% \text{ Inhibition} = 100 - \frac{\text{net stimulated release in presence of drug} \times 100}{\text{net stimulated release in absence of drug}}$$

Compounds of formula I were tested at six concentrations (100, 50, 20, 10, 5 and 1 μM) and IC_{50} values calculated. Compounds with IC_{50} values < 100 μM are considered to be active. Advantageous compounds have an IC_{50} value < 50 μM .

Medium A (for peritoneal lavage)

To a sterile 100 ml measuring cylinder was added:-
40 ml TC199 with Earle's salts (tenfold concentrate) (ICN); 4 ml heat inactivated swine serum (ICN); 10 ml sodium bicarbonate (7.5% in sterile water); 0.4 ml antibiotics solution (60 mg/ml benzylpenicillin +

- 47 -

100 mg/ml streptomycin) and 0.72 ml heparin (5000 U/ml). This mixture was transferred to sterile flask and made up to 400 ml with sterile water.

Medium B (for cell culture)

- 5 To a sterile 250 ml measuring cylinder was added:-
65 ml TC 199 (tenfold concentrate) with Earle's salts (ICN); 6.5 ml heat inactivated swine serum; 16.25 ml sodium bicarbonate (7.5% in sterile water); 0.65 ml antibiotics solution as above and 65 mg glutamine. This
10 mixture was transferred to a sterile beaker and made up to 650 ml with sterile water.

Medium C = medium B + stimulant (zymosan)

- The zymosan stimulant was prepared as follows:-
zymosan (200mg) (supplied by Sigma) was added to PBS
15 (20 ml). The mixture was boiled for 30 minutes and the volume restored to 20ml with water. The zymosan was harvested by centrifugation at 500Xg for 5 minutes, washed twice by resuspension in PBS (10ml) and centrifugation. After the final separation, the zymosan
20 was resuspended in 20ml PBS and stored as 1ml aliquots at -20°C.

650 ml medium B containing 15 ml zymosan = 12.5 particles per cell was made up and then stored in 3 ml aliquots in freezer.

- 25 Test B was carried out in the following way:

Carrageenan-induced rat paw oedema test

Female rats, weight range 125-150 g were fasted overnight. One of the hind legs of each animal was marked with a line at the connection between the cuboid/navicular and calcaneus/talus bones. Groups of six rats were orally dosed at 10 ml/kg, in random order, with a given dose of the test compound given as a solution or suspension in 10% (w/v) aqueous acacia solution.

One hour after dosing, 0.1 ml of 1% (w/v) sterile carrageenan λ in normal saline was injected deeply into the plantar surface of the marked hind foot of each rat. The volume of the foot (up to the marked line) was measured immediately after injection using duplicate water displacement readings. Three hours after injection the foot volume was measured again and the percentage increase in foot volume relative to the initial reading was calculated.

The increase in foot volume (i.e. the degree of oedema) in drug treated animals when compared with that in the drug untreated control gave the degree of inhibition of paw oedema by the drug.

Compounds were considered to be active in this test if they produced a statistically significant (typically 20% or greater) inhibition of paw oedema in at least two out of three tests after oral dosing at 100 mg/kg. Statistical significance was assessed using the Student's t test for single dose studies and Dunnett's test for multiple dose studies. More advantageous compounds were active in both Tests A and B,

TABLE 1

	Example	Test A IC ₅₀ μ M	Test B % inhibition at 100 mg/kg
	1 & 2	38	27
	3	6	
5	4	27	15
	5	35*	2
	6		
	7	11	23
	8	9	34
10	9	9	63
	10	12	47
	11	23	54
	12	7	52
	13	15	46
15	14	36	23
	15	14	31
	16	13	56
	17	15	39
	18 & 24	8*	54
20	19	10	62
	20	16	19
	21	11	
	22	28	38
	23	25	11
25	25	12	
	26	96	
	27	27	9
	28	22*	46
	29	22	13
30	30	14	10
	31	3	15

Example	Test A IC ₅₀ μ M	Test B % inhibition at 100 mg/kg
32	2	7
33	40	33
34	28	36
35	13	32
36	28	21
37	17	31
38	30	
39	1*	21
43	14*	
44	15	26
45	16	15
46	15	40

* IC₅₀ μ M based on 3 concentrations only namely 100, 10 and 1 μ M.

15 TEST C

The most advantageous compounds of formula I were active in Tests A and B and also in the following test. Carrageenan-induced pleurisy in rats was carried out as described by Ackerman et al. J. Pharmacol. Exp. Therap. 1980, 215, 588-595. Migrating leukocytes were harvested by lavage of the thoracic cavity 72h after injection of 0.3 ml 1% λ carrageenan in sterile isotonic saline. Test compounds were administered p.o. at the time of challenge and 24h and 48h thereafter.

For example the final products of Examples 7, 9, 11, 13, 16, 18 and 22 were active at 30 mg/kg in Test C and the final products of Examples 8, 10, 12 and 20 were active at 10 mg/kg or less in Test C.

Compounds which exhibit activity in both Tests A and C possess an advantageous profile of pharmacological activity unknown in commercially available antiinflammatory drugs and it is highly likely that such compounds possess significant advantages in terms of increased efficacy and/or reduced side-effects compared with known commercially available antiinflammatory drugs.

10 TEST D

Compounds of formula I suitable for the treatment of asthma were active in the three tests above and also in the late phase of the following test. Early and late phase bronchoconstriction in guinea-pigs following antigen challenge was determined by a variation of the method described by Hutson et al. Am. Rev. Respir. Dis. 1988, 137, 548-557. Guinea-pigs were sensitised by a single i.p. injection of 10 µg ovalbumin and challenged 15 to 17 days later by exposure to aerosolized antigen (4%) for five minutes, following pretreatment with mepyramine to prevent anaphylaxis. Changes in lung function were determined by whole body plethysmography at various times after challenge. Test compounds were administered p.o. 24h and 2h prior to challenge.

TEST E

The therapeutic activity of compounds of the present invention has also been demonstrated by an in vitro mixed lymphocyte reaction which is an in vitro correlate of in vivo cellular immune reactivity. A mixed lymphocyte reaction (MLR) occurs when lymphocytes

from two genetically dissimilar individuals, or inbred strains of mice are cultured together. An MLR results in the activation and proliferation of lymphocytes, which is measured by the incorporation of radiolabelled thymidine into the cellular DNA synthesised during cell division. A 'one-way' MLR is used to determine the immunosuppressive activity of test compounds in vitro. In this reaction one population of spleen cells serves as the stimulator cells and is treated with mitomycin C to prevent cell division. Spleen cells from a second allogeneic population (responder cells) are untreated and when mixed with the stimulator cells are able to undergo division which is measured. The degree of proliferation is measured in the presence and absence of test compound to assess the immunosuppressive activity of the compound.

The techniques for carrying out lymphocyte proliferation assays including MLRs are well known, eg Bradley, pages 156-166, in Mishell and Shiigi, Eds. Selected Methods in Cellular Immunology (Freeman, San Francisco, 1980); and Battisto et al, methods in Enzymology 1987; 150: 83-91. Various slight modifications of these techniques are in use and that used herein is described in Gibb, Webber and Bowen, J Immunological Methods 1985; 81: 107-113. The immunomodulant activity of compounds was determined in a mixed lymphocyte reaction in the following manner:-

Mixed Lymphocyte Reaction

Cell suspensions obtained from the spleens of female BALB/c and C57BL/6 strain mice of between 6 and 9 weeks of age were used as sources of responder and stimulator cells respectively. The mice were killed

using a rising concentration of CO₂ and the spleens removed aseptically and teased using a scalpel and forceps to produce a single cell suspension in Hanks balanced salt solution (HBSS). The suspensions were
5 filtered through cell strainers (Falcon), sedimented by centrifugation and resuspended in Tris-buffered ammonium chloride pH 7.2 (medium D) to lyse the erythrocytes. The cells were sedimented again and washed twice in HBSS before resuspending in complete RPMI 1640 tissue culture
10 medium (medium E). The C57BL/6 cells were resuspended to 9 mls per spleen and a solution of mitomycin C at 400 µg/ml in medium E added to give a final concentration of 40 µg/ml. After incubation of the C57BL/6 cells at 37°C for 30 minutes in an atmosphere of
15 5% CO₂ and 95% air, the cells were sedimented and washed 3 times in medium E. Both cell suspensions were diluted in medium E to 5×10^6 cells/ml. 100 µl of each of the responder and stimulator cell suspensions were aliquoted into the wells of 96 well flat bottom microtitre plates
20 containing 50 µl of test compound at an appropriate dilution (initially 50 µM diluted in medium E from a stock solution at 100 mM in dimethyl sulphoxide) giving a final test compound concentration of 10 µM (final dilution). Compounds active at this concentration were
25 tested at further final dilution of 1 µM, 0.1 µM and 0.01 µM to determine the concentration of test compound which causes 50% inhibition of the immune response (IC₅₀). After four days of culture at 37°C in a humidified atmosphere of 5% CO₂ and 95% air, 1 µCi of
30 tritiated thymidine in 20 µl of medium E was added to each well and the plates incubated for a further 24 hours. Cells were then harvested onto Inotech G7 glass fibre filter mats using an Inotech cell harvester. The filters were transferred to vials to which 2 mls of
35 Packard Emulsifier-Safe scintillation fluid was added and the radioactivity from the incorporated thymidine

- 54 -

- was counted in a Packard liquid scintillation counter. The counts per minute (CPM) reflect the degree of lymphocyte proliferation. The data were analysed to determine the percentage inhibition of lymphocyte proliferation by the test compound. Compounds were deemed active » if they reproducibly inhibited proliferation by $\geq 50\%$ in the absence of toxicity at a concentration of $\leq 10 \mu\text{M}$.

Medium D (for erythrocyte lysis)

10 Stock solutions

0.17M Tris. Tris base (20.6 g) was dissolved in distilled water (900 ml) and the pH adjusted to 7.65 with dilute hydrochloric acid. The volume was made up to 1000 ml with distilled water.

- 15 0.16M Ammonium chloride: Ammonium chloride (8.3 g) was dissolved in distilled water (1000 ml).

Working solutions

- The Tris stock solution (10 ml) and ammonium chloride stock solution (90 ml) were mixed, the pH adjusted to 7.2 with dilute hydrochloric acid and the solution filter sterilised.

Medium E (for cell culture)

- 25 RPMI 1640 tissue culture medium containing 2.0 g/l sodium bicarbonate (ICN FLOW) supplemented with 5 - 10% serum supplement (foetal calf serum, Sigma or Nu-Serum, Collaborative Biomedical Products) 2mM L-glutamine (ICN FLOW), 50 IU/ml penicillin (ICN FLOW), 50 $\mu\text{g/ml}$

- 55 -

streptomycin (ICN FLOW) and 5×10^{-5} M 2-mercaptoethanol (SIGMA).

MLR data were expressed as the percentage inhibition of lymphocyte proliferation caused by the test compound, calculated by the formula

$$\% \text{ inhibition} = 100 - \frac{\text{test} \times 100}{\text{control}}$$

where:

control = CPM thymidine incorporation by responder cells mixed with stimulator cells in the absence of compound

test = CPM thymidine incorporation by responder cells mixed with stimulator cells in the presence of test compound.

The final products of the Examples listed below affected the MLR at a concentration of 10 μ M in at least two out of three tests. The concentration of test compound which causes 50% inhibition of the immune response (IC_{50} μ M) for each compound is given below. A precise IC_{50} value was not determined for those examples listed as having an IC_{50} of >10 .

Example	3	4	7	8	9	10	11	12
IC₅₀	2.1	>10	2.7	1.1	3.2	9.5	2.5	1.3

Example	13	15	16	17	19	20	21	23
IC₅₀	3.2	>10	63	2.8	<1	>10	>10	>10

5

Example	26	27	29	30	31	32	33	34
IC₅₀	>10	>10	>10	>10	1.7	1.5	>10	38.7

Example	35	36	43	45	46
IC₅₀	53.8	45.3	2.01	>10	>10

The invention is illustrated by the following

10 non-limitative Examples in which compositions of mixed solvents are given by volume. Novel compounds were characterised by one or more of the following: elemental analysis, nuclear magnetic resonance, infra-red and mass spectroscopy. Temperatures are given in degrees

15 Celsius. The abbreviations HPLC (high performance liquid chromatography), THF (tetrahydrofuran), DMF (dimethylformamide), DMSO (dimethyl sulphoxide), IPA (propan-2-ol), IMS (industrial methylated spirit) and Wt (Weight), have been used in the Examples.

EXAMPLE 1

N,N-Dimethylformamide dimethyl acetal (23.3 ml) was added to 4-aminomethylpiperidine (20.0 g) in toluene (20 ml) and the mixture was boiled under reflux for 5 3 hours removing the water formed using a Dean and Stark apparatus to give a solution of 4-(dimethylaminomethyl-iminomethyl)piperidine. The mixture was cooled to ambient temperature and 2-bromoethyl phenyl ether (22.1 g) then triethylamine (24.3 ml) were added. The 10 mixture was boiled under reflux for 10 hours and then evaporated to dryness. The residue was dissolved in 2M hydrochloric acid (200 ml) and heated on a steam bath for 24 hours. The mixture was cooled to ambient temperature and washed with dichloromethane. The 15 aqueous layer was basified with 6M sodium hydroxide solution and extracted with dichloromethane to give an oil which was distilled at 186-242°C at 0.8 mbar. The distillate was dissolved in methanol (50 ml) and treated with concentrated aqueous ammonia solution (20 ml, 20 specific gravity 0.88) and then left at ambient temperature for 18 hours. The solution was concentrated by evaporation and water (200 ml) was added. The mixture was extracted with dichloromethane to give an oil. The oil was dissolved in ether and treated with 25 ethereal hydrogen chloride. A solid was collected by filtration and recrystallised from propan-2-ol/IMS to give 4-aminomethyl-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 212-214°C.

EXAMPLE 2

A mixture of 4-aminomethylpiperidine (25.0 g), acetophenone (25.6 ml), toluene (1 l) and a catalytic amount of p-toluenesulphonic acid was boiled under
5 reflux for 6 hours, removing the water formed using a Dean and Stark apparatus. The mixture was cooled to approximately 70°C then molten 2-bromoethyl phenyl ether (44.1 g) and triethylamine (30.5 ml) were added. The mixture boiled under reflux for 5 hours, then cooled and
10 evaporated. The residue was heated on a steam bath for 5 hours with 6M hydrochloric acid (400 ml), to give after work up, 4-aminomethyl-1-(2-phenoxyethyl)-piperidine dihydrochloride, m.p. 212-214°C.

EXAMPLE 3

A mixture of 4-(dimethylaminomethylimino-methyl)piperidine (7.2 g, prepared as in Example 1), 2-bromoethyl 4-chlorophenyl ether (10.0 g), toluene (100 ml) and triethylamine (5.9 ml) was heated on a steam bath for 6 hours. The mixture was evaporated to
20 dryness and the residue was dissolved in a 1:1 mixture of methanol/concentrated aqueous ammonia solution (S.G. 0.880, 80 ml). The mixture was stirred at ambient temperature for 16 hours and then evaporated to dryness. Water (50 ml) was added to the residue and the mixture
25 was extracted with dichloromethane to give an oil. The oil was dissolved in ether and acidified with ethereal hydrogen chloride to give 4-aminomethyl-1-[2-(4-chlorophenoxy)ethyl]piperidine dihydrochloride m.p. 250-252°C.

EXAMPLE 4

In a similar manner to Example 3, a mixture of 4-(dimethylaminomethyliminomethyl)piperidine (7.2 g), 3-bromopropyl phenyl ether (9.13 g), triethylamine (5.9 ml) and toluene (100 ml) was heated on a steam bath for 6 hours to give 4-aminomethyl-1-(3-phenoxypropyl)-piperidine dihydrochloride, m.p. 235°C.

EXAMPLE 5

In a similar manner to Example 3, a mixture of 4-(dimethylaminomethyliminomethyl)piperidine (7.2 g), 2-bromoethyl 4-methoxyphenyl ether (9.8 g), triethylamine (5.9 g) and toluene (100 ml) was heated on a steam bath for 6 hours to give 4-aminomethyl-1-[2-(4-methoxyphenoxy)ethyl]piperidine dihydrochloride, m.p. 245-247°C.

EXAMPLE 6

In a similar manner to Example 3, a mixture of 4-(dimethylaminomethyliminomethyl)piperidine (6.42 g), 2-bromoethyl 4-fluorophenyl ether (8.3 g), triethylamine (5.3 ml) and toluene (20 ml) was heated on a steam bath for 16 hours to give 4-aminomethyl-1-[2-(4-fluorophenoxy)ethyl]piperidine as an oil.

EXAMPLE 7

A mixture of 4-aminomethylpiperidine (3.58 g) and 4'-chloroacetophenone (4.85 g) in toluene (100 ml) was boiled under reflux with a catalytic amount of

p-toluenesulphonic acid for 5 hours, removing the water formed by means of a Dean and Stark apparatus, to give a solution. The solution was cooled and 2-bromoethyl 4-methylphenyl ether (6.68 g) was added, followed by triethylamine (4.4 ml). This mixture was boiled under reflux for 6 hours then evaporated to dryness. 6M Hydrochloric acid (50 ml) was added to the residue and the mixture heated on a steam bath for 16 hours. The mixture was evaporated to near dryness, water (50 ml) was added and the mixture was washed with ether. The aqueous layer was basified with 2M sodium hydroxide solution and extracted with dichloromethane to give 4-aminomethyl-1-[2-(4-methylphenoxy)ethyl]piperidine as an oil. The oil was cooled in ice and 98% formic acid (5.0 ml) and formaldehyde (5.0 ml, 38% aqueous solution) were added. The mixture was heated on a steam bath for 4 hours then evaporated to near dryness and the residue treated with 2M hydrochloric acid (50 ml). The solution obtained was washed with dichloromethane, basified with 2M sodium hydroxide solution and extracted with dichloromethane to give an oil which was distilled at 150°C at 0.13 mbar. This distillate was dissolved in ether and the solution acidified with ethereal hydrogen chloride. The precipitate was collected by filtration, washed with ether and recrystallised from absolute ethanol to give 4-dimethylaminomethyl-1-[2-(4-methylphenoxy)ethyl]piperidine dihydrochloride, m.p. 278-280°C.

EXAMPLE 8

In a similar manner to Example 7, a mixture of 4-aminomethylpiperidine (2.9 g), 4'-chloroacetophenone (3.94 g), toluene (100 ml) and a catalytic amount of p-toluenesulphonic acid was boiled under reflux for five

hours with removal of the water formed using a Dean and Stark apparatus. The mixture was cooled and treated with 2-bromoethyl 3-chlorophenyl ether (6.0 g) followed by triethylamine (3.54 ml). The mixture was boiled under reflux for 6 hours and then evaporated to dryness. 6M Hydrochloric acid (50 ml) was added and the mixture heated on a steam bath for 16 hours to give 4-aminomethyl-1-[2-(3-chlorophenoxy)ethyl]piperidine, as an oil. The oil, obtained on work up, was treated with 98% formic acid (5.0 ml) and formaldehyde (5.0 ml, 38% aqueous solution). The mixture was heated on a steam bath for 4 hours. After extractive work up as described in Example 7, the oil obtained was distilled at 120°C at 0.13 mbar. The distillate was treated with ethereal hydrogen chloride to give 1-[2-(3-chlorophenoxy)ethyl]-4-(dimethylaminomethyl)piperidine dihydrochloride, m.p. 258-260°C, after recrystallisation from absolute ethanol.

EXAMPLE 9

A mixture of 4-aminomethylpiperidine (7.0 g), benzaldehyde (5.77 g), toluene (70 ml) and a catalytic amount of p-toluenesulphonic acid was boiled under reflux for 7 hours with water removal using a Dean and Stark apparatus. The mixture was cooled to ambient temperature. 2-Bromoethyl phenyl ether (12.34 g) and triethylamine (8.5 ml) were added and the mixture was boiled under reflux for 5 hours. The mixture was evaporated to dryness and the residue dissolved in absolute ethanol (100 ml). Sodium borohydride (4.6 g) was added and the mixture was boiled under reflux for 18 hours. The mixture was evaporated to dryness and the residue partitioned between 2M hydrochloric acid and ethyl acetate. The aqueous layer was separated and

- basified with concentrated sodium hydroxide solution. This mixture was extracted with dichloromethane to give an oil which was dissolved in ether and treated with ethereal hydrogen chloride solution. A solid was
- 5 collected by filtration and recrystallised from IMS to give 4-(N-benzylaminomethyl)-1-(2-phenoxyethyl)-piperidine dihydrochloride, m.p. 246-247°C.

EXAMPLE 10

- 4-Aminomethyl-1-(2-phenoxyethyl)piperidine
- 10 (33.33 g) was cooled in ice and treated with 98% formic acid (33 ml) followed by formaldehyde (33 ml, 38% aqueous solution). The mixture was heated on a steam bath for 4 hours then evaporated to dryness. The residue was acidified with 3M hydrochloric acid and then
- 15 washed with ether. The acid layer was basified with 2M sodium hydroxide solution and then extracted with dichloromethane to give an oil which was distilled at 150°C at 0.13 mbar. The distillate was dissolved in ether and acidified with ethereal hydrogen chloride to
- 20 give a solid which was collected by filtration, washed with ether, and recrystallised from absolute ethanol to give 4-(dimethylaminomethyl)-1-(2-phenoxyethyl)-piperidine dihydrochloride, m.p. 268-269°C.

EXAMPLE 11

- 25 In a similar manner to Example 10, a mixture of formaldehyde (4.14 ml, 37% aqueous solution), 4-(N-benzylaminomethyl)-1-(2-phenoxyethyl)piperidine (7.8 g) and 98% formic acid (4.42 ml) was heated on a steam bath for 2.5 hours to give 4-(N-benzyl-N-methylaminomethyl)-
- 30 1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 260-262°C.

EXAMPLE 12

- In a similar manner to Example 10, a mixture of 4-aminomethyl-1-[2-(4-chlorophenoxy)ethyl]-piperidine (4.32 g), 98% formic acid (3.1 ml) and formaldehyde solution (2.9 ml, 35% aqueous solution) was heated on a steam bath for 3 hours to give 1-[2-(4-chlorophenoxy)-ethyl]-4-dimethylaminomethylpiperidine dihydrochloride m.p. 259°C.

EXAMPLE 13

- In a similar manner to Example 10, a mixture of 4-aminomethyl-1-[2-(4-fluorophenoxy)ethyl]piperidine (2.69 g, see Example 6), 98% formic acid (2.5 ml) and formaldehyde (2.5 ml, 35% aqueous solution) was heated on a steam bath for 4 hours to give 4-dimethylaminomethyl-1-[2-(4-fluorophenoxy)-ethyl]piperidine dihydrochloride, m.p. 260-262°C.

EXAMPLE 14

- In a similar manner to Example 10, a mixture of 4-aminomethyl-1-[2-(4-methoxyphenoxy)ethyl]piperidine (4.35 g), 98% formic acid (3.2 ml) and formaldehyde (3.0 ml, 35% aqueous solution) was heated on a steam bath for 3 hours to give 4-dimethylaminomethyl-1-[2-(4-methoxyphenoxy)ethyl]piperidine dihydrochloride, m.p. 250°C.

EXAMPLE 15

In a similar manner to Example 10, a mixture of 4-aminomethyl-1-(3-phenoxypropyl)piperidine (3.25 g), 98% formic acid (2.5 ml) and formaldehyde (2.4 ml, 35% aqueous solution) was heated on a steam bath for 3 hours to give 4-dimethylaminomethyl-1-(3-phenoxypropyl)-piperidine dihydrochloride, m.p. 244°C.

EXAMPLE 16

A mixture of 4-(2-dimethylaminoethyl)piperidine dihydrochloride (7.0 g), 2-bromoethyl 4-methylphenyl ether (6.6 g), triethylamine (12.75 ml) and DMF (50 ml) was heated on a steam bath for 16 hours. The mixture was poured into ice/water (500 ml), basified with 2M sodium hydroxide solution and extracted with dichloromethane. The combined organic extracts were washed with water and then extracted with 2M hydrochloric acid. The combined acidic layers were basified with 2M sodium hydroxide solution and then extracted with dichloromethane to give an oil. The oil was distilled at 120°C at 0.05 mbar and the distillate was dissolved in ether, acidified with ethereal hydrogen chloride solution and filtered. The solid obtained was washed with ether and recrystallised from absolute ethanol to give 4-(2-dimethylaminoethyl)-1-[2-(4-methylphenoxy)ethyl]piperidine dihydrochloride m.p. 281-283°C.

EXAMPLE 17

In a similar manner to Example 16, a mixture of 4-(2-dimethylaminoethyl)piperidine dihydrochloride

- 65 -

(5.5 g), 2-bromoethyl 4-fluorophenyl ether (5.26 g), triethylamine (13.4 ml) and DMF (50 ml) was heated on a steam bath for 16 hours to give 4-(2-dimethylaminoethyl)-1-[2-(4-fluorophenoxy)ethyl]piperidine dihydrochloride, m.p. 260-262°C.

EXAMPLE 18

2-Bromoethyl phenyl ether (4.94 g) was added to a mixture of 4-[2-(N-benzyl-N-methylamino)ethyl]piperidine dihydrochloride (7.5 g),
10 DMF (50 ml) and triethylamine (13.7 ml). The mixture was heated on a steam bath for 18 hours and then poured into water (300 ml) and acidified with 2M hydrochloric acid. The mixture was washed with ether then basified with 2M sodium hydroxide solution and extracted with
15 dichloromethane to give a gum. The gum was dissolved in petroleum ether b.p. 60-80°C, treated with charcoal and hot filtered. The filtrate was evaporated and the residual oil was dissolved in ether, acidified with ethereal hydrogen chloride, and filtered. The solid
20 obtained was washed with ether and recrystallised from absolute ethanol to give 4-[2-(N-benzyl-N-methylamino)ethyl]-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 276-278°C.

EXAMPLE 19

25 A mixture of 4-(2-dimethylaminoethyl)piperidine dihydrochloride (7.0 g), 2-bromoethyl 3-chlorophenyl ether (7.2 g), triethylamine (12.75 ml) and DMF (50 ml) was heated on a steam bath for 16 hours then worked up as described in Example 16 to give 1-[2-(3-chlorophenoxy)ethyl]-4-(2-dimethylaminoethyl)piperidine
30 dihydrochloride, m.p. 260-264°C.

EXAMPLE 20

A mixture of 4-(2-piperidinoethyl)piperidine (3.27 g), 2-bromoethyl phenyl ether (3.35 g), triethylamine (2.3 ml) and DMF (50 ml) was heated on a steam bath for 16 hours. The mixture was worked up as described in Example 16. The crude solid hydrochloride was treated with sodium bicarbonate solution and the mixture extracted with dichloromethane to give the free base which was purified by flash chromatography on silica using methanol/triethylamine, 99:1 as the mobile phase. The oil obtained was dissolved in ether and acidified with ethereal hydrogen chloride to give 1-(2-phenoxyethyl)-4-(2-piperidinoethyl)piperidine dihydrochloride, m.p. 285°C.

15 EXAMPLE 21

A mixture of 2-bromoethyl 2-methoxyphenyl ether (7.5 g), 4-(2-dimethylaminoethyl)piperidine dihydrochloride (7.44 g), triethylamine (18.1 ml) and DMF (50 ml) was heated on a steam bath for 18 hours. The mixture was poured into ice/water (500 ml) and basified with 2M sodium hydroxide. The product was extracted into dichloromethane to give an oil. The oil was triturated with ether. The ether was decanted off and evaporated to give crude 4-(2-dimethylaminoethyl)-1-[2-(2-methoxyphenoxy)ethyl]piperidine (4.58 g) which was added to glacial acetic acid (17 ml) and 48% hydrobromic acid (57 ml). The mixture was heated on a steam bath for 4 hours and then evaporated to dryness. The residue was dissolved in water, basified with 2M sodium hydroxide solution and extracted with dichloromethane to give an oil. The oil was dissolved in ether and acidified with ethereal hydrogen chloride. The precipitate was collected by filtration and

recrystallised from propan-2-ol to give 2-{2-[4-(2-dimethylaminoethyl)piperidino]ethoxy}phenol dihydrochloride m.p. 273-275°C.

EXAMPLE 22

- 5 A mixture of 4-(2-dimethylaminoethylidene)-1-(2-phenoxyethyl)piperidine (2.0 g) (The product of Example 37 was treated with aqueous sodium bicarbonate solution and ether. The ether was separated, washed, dried and evaporated to give the free base.) in IMS
10 (200 ml) and a catalytic amount of 10% palladium on charcoal was hydrogenated at atmospheric pressure for 5 hours. The catalyst was removed by filtration. The filtrate was evaporated to give an oil which was dissolved in ether and then acidified with ethereal
15 hydrogen chloride. The precipitate was collected by filtration and recrystallised from IMS to give 4-(2-dimethylamino-ethyl)-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 286-288°C.

EXAMPLE 23

- 20 In a similar manner to Example 22, 1-(2-phenoxyethyl)-4-[2-(pyrrolidin-1-yl)ethylidene]piperidine (from Example 38) was hydrogenated to give 1-(2-phenoxyethyl)-4-[2-(pyrrolidin-1-yl)ethyl]piperidine dihydrochloride, m.p. 270°C.

EXAMPLE 24

- 30 In a similar manner to Example 22, 4-[2-(N-benzyl-N-methylamino)ethylidene]-1-(2-phenoxyethyl)-piperidine (from Example 39) was hydrogenated to give 4-[2-(N-benzyl-N-methylamino)ethyl]-1-(2-phenoxyethyl)-piperidine dihydrochloride, m.p. 274-275°C.

EXAMPLE 25

In a similar manner to Example 22, 1-[2-(4-chlorophenoxy)ethyl]-4-(2-dimethylaminoethylidene)-piperidine (Example 40) was hydrogenated to give 1-[2-(4-chloro-phenoxy)ethyl]-4-(2-dimethylaminoethyl)-piperidine dihydrochloride, m.p. 266-267°C.

EXAMPLE 26

In a similar manner to Example 22, 4-(2-dimethylaminoethylidene)-1-[2-(2-methoxyphenoxy)ethyl]piperidine (Example 41) was hydrogenated to give 4-(2-dimethylaminoethyl)-1-[2-(2-methoxyphenoxy)ethyl]piperidine dihydrochloride, m.p. 246-247°C.

EXAMPLE 27

In a similar manner to Example 22, 4-(2-dimethylaminoethylidene)-1-[2-(4-methoxyphenoxy)ethyl]-piperidine (Example 42) was hydrogenated to give 4-(2-dimethylamino-ethyl)-1-[2-(4-methoxyphenoxy)ethyl]piperidine dihydrochloride, m.p. 254-256°C.

EXAMPLE 28

n-Butyllithium (9.66 ml, 2.5M solution in hexanes) was added dropwise to a suspension of (3-dimethylaminopropyl)triphenylphosphonium bromide (11.37 g) in dry THF (70 ml) at 0°C under nitrogen with stirring. The mixture was stirred at 0°C for 30 minutes and then 1-(2-phenoxyethyl)-4-piperidone (5.25 g) in THF (20 ml) was added dropwise over 5 minutes. The mixture was stirred at 0°C for one hour, warmed to ambient temperature and then boiled under reflux for 18 hours under nitrogen. The mixture was evaporated to dryness. Dichloromethane (100 ml) was added to the residue and

- 69 -

the mixture acidified with 2M hydrochloric acid (50 ml). The mixture was separated and the organic layer extracted with dilute hydrochloric acid. The combined acidic layers were washed with dichloromethane, basified
5 with 2M sodium hydroxide solution and extracted with ether to give an oil. The oil was extracted with hot petroleum ether b.p. 60-80°C. The petroleum ether was decanted off and evaporated to yield an oil (2.45 g) which was dissolved in IMS (100 ml) and hydrogenated at
10 1 atmosphere over 10% palladium charcoal (catalytic amount). The catalyst was removed by filtration and the filtrate evaporated to give an oil which was purified by flash chromatography on silica, using methanol and then methanol/triethylamine (99:1) as the mobile phase, to
15 give the product as an oil. The oil was dissolved in ether, acidified with ethereal hydrogen chloride and filtered to give 4-(3-dimethylaminopropyl)-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 226-227°C.

EXAMPLE 29

20 In a similar manner to Example 28, n-butyllithium (25.2 ml, 2.5 M solution in hexanes) was added to a suspension of (2-dimethylaminoethyl)-triphenylphosphonium bromide (28.7 g) in dry THF (200 ml). This mixture was then treated with a solution
25 of 1-(3-phenoxypropyl)-4-piperidone (14.7 g) in THF (20 ml). The oil obtained after work up (6.24 g) was hydrogenated in IMS at 1 atmosphere over 10% palladium charcoal over 16 hours. The catalyst was removed by filtration. The filtrate was evaporated and the residue
30 was dissolved in petroleum ether, b.p. 60-80°C. The solution was dried and evaporated to give an oil which was dissolved in ether and acidified with ethereal hydrogen chloride to yield a precipitate. The precipitate was collected by filtration and

recrystallised from IMS to give 4-(2-dimethylaminoethyl)-1-(3-phenoxypropyl)piperidine dihydrochloride, m.p. 294-295°C.

EXAMPLE 30

5 In a similar manner to Example 28, a mixture of n-butyllithium (20 ml, 2.5M solution in hexanes) was added to a suspension of (2-dimethylaminoethyl)-triphenylphosphonium bromide (22.8 g) in THF (200 ml). The mixture was treated with 1-(4-phenoxybutyl)-4-
10 piperidone (12.35 g) in dry THF (20 ml). The oil obtained on work up was hydrogenated in IMS at 1 atmosphere over 10% palladium charcoal over 16 hours to give 4-(2-dimethylaminoethyl)-1-(4-phenoxybutyl)-piperidine dihydrochloride, m.p. 268-270°C.

15 EXAMPLE 31

A mixture of 4-aminomethylpiperidine (5.0 g), 4'-chloroacetophenone (6.77 g), p-toluenesulphonic acid (catalytic amount) and toluene (50 ml) was boiled under reflux for 8 hours while removing the water formed using
20 a Dean and Stark apparatus. The mixture was cooled to ambient temperature and 2-bromoethyl phenyl ether (8.84 g) and triethylamine (6.1 ml) were added. The mixture was boiled under reflux for 2 hours and evaporated to dryness. The residue was dissolved in
25 ethanol (100 ml) and sodium borohydride (3.4 g) was added. The mixture was boiled under reflux for 18 hours and then evaporated to dryness. The residue was dissolved in 2M hydrochloric acid, washed with ethyl acetate, basified with 5M sodium hydroxide solution and
30 the mixture extracted with dichloromethane to give an oil. The oil was dissolved in ether and treated with ethereal hydrogen chloride solution to give a solid

- 71 -

which was collected by filtration and recrystallised from absolute ethanol to give 4-[1-(4-chlorophenyl)-ethylaminomethyl]-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 220°C.

5 EXAMPLE 32

In a similar manner to Example 31, a mixture of 4-(2-aminoethyl)piperidine (5.0 g), 4'-chloro-acetophenone (6.04 g), p-toluenesulphonic acid (catalytic amount) and toluene (100 ml) was boiled under
10 reflux for 10 hours with water removal using a Dean and Stark apparatus. The mixture was cooled to ambient temperature and 2-bromoethyl phenyl ether (7.85 g) and triethylamine (5.4 ml) added. The mixture was boiled under reflux for 6 hours. The solvent was evaporated
15 off and the residue dissolved in ethanol (100 ml) and sodium borohydride (3.4 g) was added. The mixture was boiled under reflux for 4 hours then worked up to give an oil which was purified by flash column chromatography on silica using ethyl acetate/triethylamine (95:5) as
20 the mobile phase to give 4-{2-[1-(4-chlorophenyl)ethyl-amino]ethyl}-1-(2-phenoxyethyl)piperidine as an oil.

EXAMPLE 33

a) Acetic anhydride (8.72 g) was added with stirring to a mixture of 4-aminomethyl-1-(2-
25 phenoxyethyl)piperidine (20.0 g) in pyridine (85 ml) containing 4-dimethylaminopyridine (catalytic amount) with cooling to maintain ambient temperature. The mixture was stirred overnight at ambient temperature and then evaporated to dryness. Water was removed by adding
30 toluene and removing the toluene by distillation. The residue was dissolved in water (100 ml) and 2M hydrochloric acid (50 ml). The solution was washed with

dichloromethane, basified with concentrated sodium hydroxide solution and extracted with dichloromethane to give N-[1-(2-phenoxyethyl)piperid-4-ylmethyl]acetamide as an oil which solidified on standing, m.p. 79-80°C.

- 5 b) A solution of the acetamide from part a) (6.05 g) in dry THF (50 ml) was added dropwise with stirring to a suspension of sodium hydride (0.88 g, 60% dispersion in mineral oil) in dry THF (40 ml) with cooling at ambient temperature under nitrogen. The suspension was stirred for 30 minutes and then a solution of ethyl iodide (3.42 g) in dry THF (20 ml) was added dropwise with stirring. The mixture was stirred at ambient temperature for 5 hours and then heated on a steam bath with stirring for 18 hours. The mixture was evaporated to dryness. The residue was dissolved in 2M hydrochloric acid and washed with ethyl acetate. The aqueous layer was basified with concentrated sodium hydroxide solution and then extracted with dichloromethane to give an oil. The oil was extracted with boiling petroleum ether b.p. 60-80°C and then the extracts were combined and evaporated to dryness. The residue was stirred with petroleum ether, b.p. 60-80°C (100 ml). The solution was decanted from the residue, dried and evaporated to give N-ethyl-N-[1-(2-phenoxyethyl)piperid-4-ylmethyl]acetamide.

- c) Lithium aluminium hydride solution (11.5 ml, 1M solution in THF) was added to dry THF (11 ml) with stirring at 0°C under nitrogen. The solution was stirred for 5 minutes and then concentrated sulphuric acid (0.31 ml) was added dropwise at 0°C. The mixture was stirred for 15 minutes then a solution of the product from b) (2.65 g) in dry THF (15 ml) was added dropwise at 0°C. The mixture was allowed to warm up to ambient temperature with stirring and then stirred at

ambient temperature for 3 hours. The mixture was cooled in ice and a mixture of water and THF (1:1) (2 ml) was added dropwise, followed by 2M sodium hydroxide solution (7 ml). The mixture was filtered and evaporated to dryness. The residue was basified with 2M sodium hydroxide solution and extracted with dichloromethane to give an oil. The oil was dissolved in ether and treated with ethereal hydrogen chloride solution to give a solid which was collected by filtration and recrystallised from propan-2-ol to give 4-diethylaminomethyl-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 195-197°C.

EXAMPLE 34

In a similar manner to Example 33 c), *N*-[1-(2-phenoxyethyl)piperid-4-ylmethyl]acetamide (13.0 g) was reduced to give 4-(ethylaminomethyl)-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 268-270°C.

EXAMPLE 35

A solution of 4-(ethylaminomethyl)-1-(2-phenoxyethyl)piperidine (3.0 g) in 98% formic acid (2.2 ml) and 37% aqueous formaldehyde (2.06 ml) was heated on a steam bath for 5 hours. Work up as described in Example 10 gave an oil which was dissolved in ether and treated with ethereal hydrogen chloride solution to give a solid which was collected by filtration and recrystallised from absolute ethanol to give 4-(*N*-ethyl-*N*-methylaminomethyl)-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 248-250°C.

EXAMPLE 36

a) A solution of propionyl chloride (2.37 g) in dichloromethane (10 ml) was added dropwise to a solution

- 74 -

of 4-aminomethyl-1-(2-phenoxyethyl)piperidine (6.0 g) in dichloromethane (50 ml) and triethylamine (3.9 ml) at 0°C. The solution was allowed to warm up to ambient temperature with stirring and then stirred at this temperature for a further 2 hours. More propionyl chloride (0.3 ml) was added at ambient temperature and the solution stirred for a further half-hour. Further propionyl chloride (0.3 ml) was added and the mixture stirred for 1 hour. The mixture was extracted with 2M hydrochloric acid. The combined acidic extracts were washed with ethyl acetate, then basified with sodium hydroxide solution and extracted with dichloromethane to give an oil which solidified on standing to give N-[1-(2-phenoxyethyl)piperid-4-ylmethyl]propionamide, m.p. 85-86°C.

b) The product from part a) was reduced in a similar manner to Example 33 c) to give 1-(2-phenoxyethyl)-4-(propylaminomethyl)piperidine dihydrochloride, m.p. 265-267°C.

20 EXAMPLE 37

n-Butyllithium (9.13 ml, 2.5M solution in hexanes) was added to a suspension of (2-dimethylaminoethyl)triphenylphosphonium bromide (9.46 g) in dry THF (60 ml) under nitrogen at 0°C over 5 minutes. The mixture was stirred at 0°C for 30 minutes and then 1-(2-phenoxyethyl)-4-piperidone (5.0 g) in THF (15 ml) was added dropwise over 5 minutes. The mixture was warmed to ambient temperature and then heated at 60°C for 5 hours. The mixture was left at ambient temperature for 18 hours and then quenched with water and diluted with ethyl acetate. 2M Hydrochloric acid was added and the mixture was separated. The organic layer was extracted with 2M

hydrochloric acid and the combined acidic layers were basified and extracted with dichloromethane to give an oil. The oil was purified by flash column chromatography on silica using dichloromethane and then
5 dichloromethane/methanol (1:1) as the mobile phase. The oil obtained after chromatography was dissolved in ether and treated with ethereal hydrogen chloride. The ether was decanted off and the residual semi-solid was triturated with acetone/ether (1:1) and filtered to give
10 a solid which was recrystallised from IMS to give 4-(2-dimethylaminoethylidene)-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 234-235°C.

EXAMPLES 38-42

Examples 38-42 were carried out in a similar
15 manner to Example 37 as summarised in Table 1 below. A compound of formula IX, in which ALK^1 represents $-(CH_2)_2-$, Y represents a piperidine ring which has ALK^1 attached at the 1-position and the oxygen attached at the 4-position and R_1 , R_2 and R_3 are as given in Table
20 1, was reacted with a compound of formula Xa in which R_4 represents hydrogen, ALK^2 represents methylene, X represents bromide and R_5 and R_6 are as given in Table 1.

TABLE 1

Example	IX R ₁ R ₂ R ₃	Wt of IX (g)	THF (ml)	Xa		Wt of Xa (g)	n-BuLi* ml	Notes
				R ₅	R ₆			
38	H	3.6	55	-(CH ₂) ₄ -		7.2	6.5	
39	H	3.3	60	CH ₃	PhCH ₂	7.3	6.0	
40	4-Cl	5.0	80	CH ₃	CH ₃	9.0	7.9	(1)
41	2-OCH ₃	5.0	80	CH ₃	CH ₃	9.1	8.0	(1)
42	4-OCH ₃	3.0	45	CH ₃	CH ₃	5.5	4.8	(1)

5

10 (1) The acidic extracts were basified with dilute sodium hydroxide solution and extracted into ether to give the product in free base form, as an oil.

* 2.5M solution in hexanes.

- 77 -

The compounds prepared in Table 1 were as follows:

EXAMPLE 38

1-(2-Phenoxyethyl)-4-[2-(pyrrolidin-1-yl)ethylidene]
5 piperidine dihydrochloride hemihydrate, m.p. 227-228°C.

EXAMPLE 39

4-[2-(N-Benzyl-N-methylamino)ethylidene]-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 225-226°C.

EXAMPLE 40

10 1-[2-(4-Chlorophenoxy)ethyl]-4-(2-dimethylaminoethylidene)piperidine as an oil.

EXAMPLE 41

4-(2-Dimethylaminoethylidene)-1-[2-(2-methoxyphenoxy)ethyl]piperidine as an oil.

15 EXAMPLE 42

4-(2-Dimethylaminoethylidene)-1-[2-(4-methoxyphenoxy)ethyl]piperidine as an oil.

EXAMPLE 43

20 n-Butyllithium (4.1 ml, 2.5M solution in hexanes) was added to a mixture of 2-dimethylaminoethyl triphenylphosphonium bromide (4.71 g) in dry THF (50 ml) with stirring at 0°C under nitrogen. The mixture was stirred at 0°C for 30 minutes and then a solution of N-(2-phenoxyethyl)piperidine-3-carbaldehyde (2.41 g) in

- 78 -

THF (5 ml) was added dropwise at 0°C with stirring. The mixture was allowed to warm to ambient temperature and then boiled under reflux for 18 hours under nitrogen. The mixture was evaporated to dryness under reduced
5 pressure and then treated with dilute hydrochloric acid. The mixture was washed with dichloromethane, basified with 2M sodium hydroxide solution and extracted with dichloromethane to give an oil. The oil was triturated with petroleum ether b.p. 40-60°C, and the supernatant
10 petroleum ether decanted off to leave an oil. The oil was stored at 5°C for 64 hours, during which time a fine crystalline solid formed. Petroleum ether b.p. 40-60°C was added and the mixture was filtered to give a solid which was discarded. The filtrate was evaporated to
15 yield a pale yellow oil, which was identified as (E)-3-(3-dimethylamino-1-propenyl)-1-(2-phenoxyethyl)piperidine. A portion of this oil (2.0 g) was dissolved in ether and acidified with ethereal citric acid to give a precipitate which was collected by
20 filtration, washed with acetone and dried to give (E)-3-(3-dimethylamino-1-propenyl)-1-(2-phenoxyethyl)-piperidine dicitrate, m.p. 65-70°C.

EXAMPLE 44

(E)-3-(3-dimethylamino-1-propenyl)-1-(2-
25 phenoxyethyl)piperidine (5.0 g) was hydrogenated in a similar manner to Example 22 to give 3-(3-dimethylamino-propyl)-1-(2-phenoxyethyl)piperidine dihydrochloride, m.p. 204-206°C (with decomposition).

EXAMPLE 45

30 A mixture of 3-chloromethyl-1-(2-phenoxyethyl)piperidine hydrochloride (4.2 g), pyrrolidine (1.23 g), sodium bicarbonate (3.7 g),

- 79 -

- potassium iodide (catalytic amount) and IMS (100 ml) was boiled under reflux for 48 hours. The mixture was evaporated to dryness under reduced pressure and ether was added to the residue. The mixture was filtered.
- 5 The filtrate was evaporated under reduced pressure to give a residue to which pyrrolidine (30 ml) was added. This mixture was boiled under reflux for 16 hours. The excess pyrrolidine was removed under reduced pressure and the residue was purified by azeotropic distillation
- 10 with toluene. The residue was partitioned between ether and water and the ether layer was separated. The aqueous layer was extracted with ether. The combined ether extracts yielded an oil which was dissolved in ether and acidified with ethereal citric acid to give a
- 15 solid which was collected by filtration to give 1-(2-phenoxyethyl)-3-(pyrrolidin-1-ylmethyl)piperidine dicitrate, m.p. 69°C (with decomposition).

EXAMPLE 46

- A mixture of 3-chloromethyl-1-(2-phenoxyethyl)piperidine (2.8 g) and piperidine (15 ml)
- 20 was heated on a steam bath for 24 hours. The excess piperidine was removed under reduced pressure and the residue purified by azeotropic distillation with toluene. The residue was dissolved in ether/water and
- 25 then basified to pH 14. The mixture was extracted with ether and the combined ether extracts yielded an oil which was dissolved in ether and acidified with ethereal citric acid to give a solid which was collected by filtration to give 1-(2-phenoxyethyl)-3-(piperidin-1-
- 30 ylmethyl)piperidine dicitrate, m.p. 88-89°C (with decomposition).

PREPARATION OF STARTING MATERIALS

Starting materials, not detailed below, were commercially available and may be found by reference to the Fine Chemicals Directory or suppliers' catalogues.

5 [2-(Pyrrolidin-1-yl)ethyl]triphenylphosphonium bromide

a) A mixture of 2-bromoethyl phenyl ether (20.0 g), triphenylphosphine (26.2 g) and phenol (200 g) was heated at 90°C with stirring for 48 hours. The mixture was cooled to approximately 40°C and added to
10 ether (1500 ml). The ether was decanted off and the residual oil was triturated with ether until a solid was obtained. The solid was collected by filtration to give (2-phenoxyethyl)triphenylphosphonium bromide.

b) A mixture of the product from a) (10 g),
15 pyrrolidine (5.0 g) and DMSO (20 ml) was heated at 40°C for 10 minutes with stirring. The mixture was poured into ether. The ether was decanted off and the residual oil was triturated with ether to give a solid which was collected by filtration to give
20 [2-(pyrrolidin-1-yl)ethyl]triphenylphosphonium bromide, m.p. 188-190°C.

The other phosphonium salts required as starting materials in Table 1 were prepared in a similar manner, by reacting (2-phenoxyethyl)triphenylphosphonium bromide with the appropriate amine.

25 4-[2-(Chlorophenoxy)ethyl]piperidone

A mixture of 2-bromoethyl 4-chlorophenyl ether (10.0 g), 4-piperidone (6.5 g), sodium bicarbonate (7.14 g) and IMS (100 ml) was boiled under reflux for 24 hours. The mixture was cooled and the solvent

- 81 -

removed under reduced pressure. Water was added to the residue and the mixture was extracted with dichloromethane. The combined organic extracts were then extracted with dilute hydrochloric acid and the
5 acidic extracts were filtered and then basified with sodium hydroxide solution. The mixture obtained was extracted with ether to give 4-[2-(chlorophenoxy)-ethyl]piperidone as a solid.

The other 1-substituted 4-piperidones required
10 as starting materials in Table 1 were prepared in a similar manner by reacting 4-piperidone with the appropriately substituted commercially available halo ether.

(3-Dimethylaminopropyl)triphenylphosphonium bromide

15 a) 1,3-Dibromopropane (28.8 ml) was added to a solution of triphenylphosphine (15.0 g) in toluene (30 ml). The mixture was stirred at 70°C under nitrogen for 16 hours, then cooled, diluted with ether (150 ml) and filtered to give 3-bromopropyltriphenylphosphonium
20 bromide.

b) A solution of dimethylamine in IMS (80 ml of a 33% solution) was added to a suspension of the product from a) (12.5 g) in absolute ethanol (160 ml) at 0°C. The mixture was stirred at ambient temperature for 5
25 hours and then left to stand at this temperature for 18 hours. The solvent was removed under reduced pressure and the residue was triturated with dichloromethane. The mixture was filtered to give (3-dimethylaminopropyl)triphenylphosphonium bromide,
30 m.p. 275°C.

2-Chloroethyl 4-chlorophenyl ether

A mixture of 4-chlorophenol (100 g), 1-bromo-2-chloroethane (335 g), potassium carbonate (129 g) and acetone (2 l) was boiled under reflux with stirring for 18 hours. The mixture was evaporated to dryness under reduced pressure and then dichloromethane was added to the cool residue. The mixture was filtered and the filtrate washed with 5M sodium hydroxide solution, water, dried, filtered and evaporated to give 2-chloroethyl 4-chlorophenyl ether as an oil.

4-(2-Dimethylaminoethyl)piperidine

a) Benzoyl chloride (25.8 ml) in dichloromethane (20 ml) was added dropwise over 30 minutes to a mixture of 4-(2-hydroxyethyl)piperidine (26 g) in dichloromethane (250 ml) and triethylamine (16.3 ml) at 0°C. The mixture was allowed to warm to ambient temperature and stirred at this temperature for 1 hour. The mixture was washed with sodium bicarbonate solution and then dried and evaporated to give a colourless oil. The oil was dissolved in 1,4-dioxane (250 ml) and then a solution of sodium hydroxide (5.0 g) in water (20 ml) was added and the mixture stirred at ambient temperature for 16 hours and then at 60°C for 6 hours. The mixture was evaporated to near dryness, basified with 2M sodium hydroxide solution and the product extracted into dichloromethane to give 1-benzoyl-4-(2-hydroxyethyl)-piperidine as an oil.

b) Thionyl chloride (0.38 ml) was added dropwise to a solution of the product from a) (1.0 g) in chloroform (10 ml) with stirring. The mixture was stirred at ambient temperature for 1 hour and then boiled under reflux for 30 minutes. The mixture was

evaporated to dryness and the residue was dissolved in dichloromethane, washed with sodium bicarbonate solution, dried and evaporated to give 1-benzoyl-4-(2-chloroethyl)piperidine as an oil.

- 5 c) A mixture of the product from b) (11.61 g) and a 33% solution of dimethylamine in absolute ethanol (100 ml) was boiled under reflux for 16 hours. The mixture was evaporated to dryness and 2M hydrochloric acid (50 ml) was added. The mixture was washed with
10 ether and then the acidic layer was basified with 2M sodium hydroxide solution and the product extracted into dichloromethane to give a gum. The gum was dissolved in petroleum ether b.p. 60-80°C and treated with charcoal, then filtered and the filtrate evaporated to give an
15 oil. A portion of the oil (1.5 g) was dissolved in ether and acidified with ethereal oxalic acid to give 1-benzoyl-4-(2-dimethylaminoethyl)piperidine oxalate, m.p. 82-84°C.

- d) The oil from c) (6.0 g) and 6M hydrochloric acid (50 ml) was heated on a steam bath for 16 hours. The mixture was evaporated to near dryness and basified with 2M sodium hydroxide solution. The mixture was extracted with dichloromethane to give an oil which was dissolved in ether and acidified with ethereal hydrogen
20 chloride. The mixture was filtered to give a solid which was recrystallised from IMS/IPA (1:1) to give 4-(2-dimethylaminoethyl)piperidine dihydrochloride, m.p. 242-245°C.

4-(2-Piperidinoethyl)piperidine

- 30 a) A mixture of 1-benzoyl-4-(2-chloroethyl)piperidine (5.86 g), piperidine (11.52 ml) and IMS (100 ml) was boiled under reflux for 10 hours.

- The mixture was evaporated to near dryness and then dried by azeotropic distillation with toluene. The residue was dissolved in dichloromethane, washed with 2M sodium hydroxide solution and then brine. The separated organic layer was extracted with 2M hydrochloric acid and the combined acidic layers were basified with 2M sodium hydroxide solution and then extracted with dichloromethane to give 1-benzoyl-4-(2-piperidinoethyl)piperidine as an oil.
- 10 b) The product from a) (5.81 g) and 6M hydrochloric acid (100 ml) were heated on a steam bath for 16 hours. The mixture was evaporated to dryness then diluted with water (100 ml) and then washed with dichloromethane. The aqueous layer was basified with 2M sodium hydroxide solution and then extracted with dichloromethane to give 4-(2-piperidinoethyl)piperidine as an oil.

4-[2-(N-Benzyl-N-methylamino)ethyl]piperidine

- 20 a) A mixture of 1-benzoyl-4-(2-chloroethyl)piperidine (11.61 g), N-methylbenzylamine (16.75 g) and IMS (100 ml) was boiled under reflux for 16 hours. The mixture was evaporated to dryness and then 4M hydrochloric acid (50 ml) was added to the residue. The mixture was washed with ether and the acidic layer was then basified with 4M sodium hydroxide solution and extracted with dichloromethane to give an oil. Excess amine was distilled off at 13 mbar and the residue was triturated with petroleum ether b.p. 40-60°C. The supernatant petroleum ether was decanted off and the residual oil was dried under vacuum to give 1-benzoyl-4-[2-(N-benzyl-N-methylamino)ethyl]piperidine. A small portion of the oil was dissolved in ether and acidified with ethereal oxalic acid. The supernatant

liquid was decanted off and the solid residue was triturated with acetone and filtered to give the oxalate salt, m.p. 141-142°C.

- b) The oil from a) (5.0 g) and 6M hydrochloric acid (50 ml) were heated together on a steam bath for 16 hours. The mixture was evaporated to near dryness and the residue basified with 2M sodium hydroxide solution. The mixture was extracted with dichloromethane to give an oil which was dissolved in ether, acidified with ethereal oxalic acid and then filtered to give a solid which was recrystallised from IMS to give 4-[2-(N-benzyl-N-methylamino)ethyl]-piperidine dioxalate, m.p. 95-96°C.

N-(2-Phenoxyethyl)piperidine-3-carbaldehyde

- a) A mixture of ethyl 3-piperidinecarboxylate (20.58 g), 2-bromoethyl phenyl ether (26.31 g), sodium bicarbonate (10.99 g) and IMS (100 ml) was boiled under reflux for 16 hours. The mixture was evaporated to dryness then diluted with water (100 ml) and 2M sodium hydroxide solution (20 ml). This mixture was extracted with dichloromethane and the combined dichloromethane extracts were extracted with 2M hydrochloric acid. The acidic extracts were washed with dichloromethane and then basified with 2M sodium hydroxide solution. This mixture was extracted with dichloromethane to give ethyl N-(2-phenoxyethyl)piperidine-3-carboxylate as an oil. A sample of the oil was dissolved in ether and ethereal oxalic acid was added. The precipitate was filtered off and recrystallised from aqueous propan-2-ol to give ethyl N-(2-phenoxyethyl)piperidine-3-carboxylate oxalate hemihydrate, m.p. 131-132°C.

- 86 -

b) Di-isobutylaluminium hydride (13.2 ml of 1.5M solution in toluene) was added to a mixture of ethyl N-(2-phenoxyethyl)piperidine-3-carboxylate (5.0 g) in dry THF (50 ml) at 0°C under nitrogen with stirring. The mixture was quenched with water (5 ml) and then treated with saturated Rochelles salt solution. The mixture was stirred at room temperature for 30 minutes and then filtered. The filtrate was separated and the organic layer was washed, dried and evaporated to give N-(2-phenoxyethyl)piperidine-3-carbaldehyde as an oil.

3-Chloromethyl-1-(2-phenoxyethyl)piperidine hydrochloride

a) A mixture of 2-bromoethyl phenyl ether (20.1 g), piperidin-3-yl-methanol (11.5 g), sodium bicarbonate (8.4 g) and IMS (100 ml) was boiled under reflux for 16 hours. The mixture was evaporated to dryness under reduced pressure and the residue was treated with water (50 ml), 2M sodium hydroxide solution (20 ml) and extracted with dichloromethane. The combined dichloromethane extracts were washed with brine and then extracted with 2M hydrochloric acid. The combined acidic extracts were washed with dichloromethane then basified with 2M sodium hydroxide solution and then extracted with dichloromethane to give 1-(2-phenoxyethyl)piperidin-3-ylmethanol as an oil.

b) A solution of thionyl chloride (1.7 ml) in chloroform (10 ml) was added dropwise with stirring to a solution of 1-(2-phenoxyethyl)piperidin-3-ylmethanol (5.0 g) in chloroform (70 ml) at 0-5°C. After the addition the mixture was boiled under reflux for 2.5 hours. The solvent was removed under reduced pressure and the residue was triturated with ether and

- 87 -

filtered to give 3-chloromethyl-1-(2-phenoxyethyl)-piperidine hydrochloride, m.p. 125-127°C.

PHARMACEUTICAL EXAMPLES

Example U

- 5 In the preparation of capsules, 10 parts by weight of active compound and 240 parts by weight of lactose are de-aggregated and blended. The mixture is filled into hard gelatin capsules, each capsule containing 10 mg active compound.

10 Example V

 Tablets are prepared from the following ingredients.

	<u>Parts by Weight</u>
Active compound	10
15 Lactose	190
Maize starch	22
Polyvinylpyrrolidone	10
Magnesium stearate	3

- 20 The active compound, the lactose and some of the starch are de-aggregated, blended and the resulting mixture is granulated with a solution of the polyvinylpyrrolidone in ethanol. The dry granulate is blended with magnesium stearate and the rest of the starch. The mixture is then compressed in a tableting machine to give tablets
- 25 containing 10 mg of active compound.

Example W

Tablets are prepared by the method of the previous Example. The tablets are enteric coated in a conventional manner using a solution of 20% cellulose acetate phthalate and 3% diethyl phthalate in ethanol:dichloromethane (1:1).

Example X

In the preparation of suppositories, 100 parts by weight of active compound is incorporated in 1300 parts by weight of semi-synthetic glycerides as the suppository base and the mixture formed into suppositories each containing 100 mg of active ingredient.

Example Y

In the preparation of capsules, 50 parts by weight of active compound, 300 parts by weight of lactose and 3 parts by weight of magnesium stearate are de-aggregated and blended. The mixture is filled into hard gelatin capsules, each capsule containing 50 mg of active ingredient.


Example Z

The active compound is incorporated into the base by thorough homogenization until the drug is evenly distributed. The ointment is packed into 10 g amber jars with screw-capped lined lids.

Active compound 0.1 g
White soft paraffin to 10 g

CLAIMS

1. Compounds of formula I



- and pharmaceutically acceptable salts thereof in which
 R_1 , R_2 and R_3 independently represent hydrogen, hydroxy,
 5 halo, a C_{1-6} alkyl group (optionally substituted by one
 or more halogen atoms) or a C_{1-6} alkoxy group
 (optionally substituted by one or more halogen atoms);

ALK^1 represents a C_{2-6} alkylene chain optionally
 substituted by one or more C_{1-2} alkyl groups;

- 10 Y represents a piperidine ring which is attached through
 nitrogen to ALK^1 ;

R_4 represents hydrogen or a C_{1-4} alkyl group;

- the broken line in --- represents a bond, so that a
 double bond connects Y and CR_4 , or is absent and the
 15 free valency on Y is taken up by hydrogen and the free
 valency on CR_4 is taken up by hydrogen or a C_{1-4} alkyl
 group;


- ALK^2 is absent or represents a C_{1-4} alkylene chain
 optionally substituted by one or more C_{1-2} alkyl groups;
 20 and

R_5 and R_6 independently represent hydrogen, a C_{1-6} alkyl
 group, a phenyl C_{1-4} alkyl group (in which the phenyl

- ring is optionally substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, halo or trifluoromethyl) or R₅ and R₆ together with the nitrogen atom to which they are attached represent a
- 5 saturated 3-7 membered heterocyclic ring, optionally containing sulphur, oxygen or an additional nitrogen atom, wherein the ring is optionally substituted by one or more C₁₋₄ alkyl groups;

- excluding 4-aminomethyl-1-[2-(4-fluorophenoxy)ethyl]-
- 10 piperidine, 4-(2-aminoethyl)-1-[2-(4-fluorophenoxy)ethyl]piperidine and 3-aminomethyl-1-[2-(2-methoxyphenoxy)ethyl]piperidine.

2. Compounds according to claim 1 as represented by formula II



- 15 and pharmaceutically acceptable salts thereof in which

R₁, R₂ and R₃ independently represent hydrogen, hydroxy, halo, trifluoromethyl, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group;

n = 2-4 and m = 1-3;


- 20 R₅ represents hydrogen, a C₁₋₄ alkyl group or a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group or halo); and

- 91 -

R₆ represents a C₁₋₄ alkyl group or a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group or halo);

- 5 or R₅ and R₆ together with the nitrogen atom to which they are attached represent a pyrrolidine ring, a piperidine ring or a morpholine ring.

3. Compounds according to either claim 1 or claim 2 as represented by formula III



- 10 and pharmaceutically acceptable salts thereof in which

R₁ represents hydrogen, halo, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group;

R₂ represents hydrogen, halo, a C₁₋₄ alkyl group or a C₁₋₄ alkoxy group;

- 15 n = 2 and m = 1-3;

R₃ represents hydrogen, a C₁₋₄ alkoxy group or hydroxy;

R₅ represents hydrogen, a C₁₋₄ alkyl group or a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally substituted by one or more of the following: a C₁₋₄

- 20 alkyl group, a C₁₋₄ alkoxy group or halo); and

R₆ represents a C₁₋₄ alkyl group or a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally

- 92 -

substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group or halo;

or R₅ and R₆ together with the nitrogen atom to which they are attached represent a pyrrolidine ring, a piperidine ring or a morpholine ring.

4. Compounds according to claim 3 in which R₁ represents hydrogen, chloro, fluoro, methoxy or methyl.

5. Compounds according to claim 3 in which R₂ represents hydrogen or chloro.

10 6. Compounds according to claim 3 in which R₃ represents hydrogen, hydroxy or methoxy.

7. Compounds according to claim 3 in which R₅ represents hydrogen, methyl, ethyl or propyl and R₆ represents methyl, ethyl, propyl or benzyl or R₅ and R₆ together with the nitrogen atom to which they are attached represent a pyrrolidine ring or a piperidine ring.

8. Compounds according to claim 3 in which m represents 1.

20 9. A compound of formula I according to claim 1 selected from:

4-(dimethylaminomethyl)-1-[2-(4-methylphenoxy)ethyl]-piperidine;


25 1-[2-(3-chlorophenoxy)ethyl]-4-(dimethylaminomethyl)-piperidine;

4-(N-benzylaminomethyl)-1-(2-phenoxyethyl)piperidine;

4-(N-benzyl-N-methylaminomethyl)-1-(2-phenoxyethyl)-piperidine;

- 4-(dimethylaminomethyl)-1-(2-phenoxyethyl)piperidine;
 1-[2-(4-chlorophenoxy)ethyl]-4-(dimethylaminomethyl)-
 piperidine;
 4-(dimethylaminomethyl)-1-[2-(4-fluorophenoxy)ethyl]-
 5 piperidine;
 4-(2-dimethylaminoethyl)-1-[2-(4-methylphenoxy)ethyl]-
 piperidine;
 4-[2-(N-benzyl-N-methylamino)ethyl]-1-(2-phenoxyethyl)-
 piperidine;
 10 1-(2-phenoxyethyl)-4-(2-piperidinoethyl)piperidine; and
 4-(2-dimethylaminoethyl)-1-(2-phenoxyethyl)piperidine.

10. A pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I



- 15 and pharmaceutically acceptable salts thereof in which R_1 , R_2 and R_3 independently represent hydrogen, hydroxy, halo, a C_{1-6} alkyl group (optionally substituted by one or more halogen atoms) or a C_{1-6} alkoxy group (optionally substituted by one or more halogen atoms);
- 20 ALK^1 represents a C_{2-6} alkylene chain optionally substituted by one or more C_{1-2} alkyl groups;
- Y represents a piperidine ring which is attached through nitrogen to ALK^1 ;
- R_4 represents hydrogen or a C_{1-4} alkyl group;


the broken line in --- represents a bond, so that a double bond connects Y and CR₄, or is absent and the free valency on Y is taken up by hydrogen and the free valency on CR₄ is taken up by hydrogen or a C₁₋₄ alkyl group;

5 group;

ALK² is absent or represents a C₁₋₄ alkylene chain optionally substituted by one or more C₁₋₂ alkyl groups; and

- R₅ and R₆ independently represent hydrogen, a C₁₋₆ alkyl group, a phenyl C₁₋₄ alkyl group (in which the phenyl ring is optionally substituted by one or more of the following: a C₁₋₄ alkyl group, a C₁₋₄ alkoxy group, halo or trifluoromethyl) or R₅ and R₆ together with the nitrogen atom to which they are attached represent a saturated 3-7 membered heterocyclic ring, optionally containing sulphur, oxygen or an additional nitrogen atom, wherein the ring is optionally substituted by one or more C₁₋₄ alkyl groups.
- 10
- 15

11. The use of compounds of formula I



- 20 and pharmaceutically acceptable salts thereof in which R₁, R₂ and R₃ independently represent hydrogen, hydroxy, halo, a C₁₋₆ alkyl group (optionally substituted by one or more halogen atoms) or a C₁₋₆ alkoxy group (optionally substituted by one or more halogen atoms);
- 25 ALK¹ represents a C₂₋₆ alkylene chain optionally substituted by one or more C₁₋₂ alkyl groups;

- 95 -

Y represents a piperidine ring which is attached through nitrogen to ALK^1 ;

R_4 represents hydrogen or a C_{1-4} alkyl group;

the broken line in --- represents a bond, so that a
5 double bond connects Y and CR_4 , or is absent and the free valency on Y is taken up by hydrogen and the free valency on CR_4 is taken up by hydrogen or a C_{1-4} alkyl group;

ALK^2 is absent or represents a C_{1-4} alkylene chain
10 optionally substituted by one or more C_{1-2} alkyl groups; and

R_5 and R_6 independently represent hydrogen, a C_{1-6} alkyl group, a phenyl C_{1-4} alkyl group (in which the phenyl ring is optionally substituted by one or more of the
15 following: a C_{1-4} alkyl group, a C_{1-4} alkoxy group, halo or trifluoromethyl) or R_5 and R_6 together with the nitrogen atom to which they are attached represent a saturated 3-7 membered heterocyclic ring, optionally containing sulphur, oxygen or an additional nitrogen
20 atom, wherein the ring is optionally substituted by one or more C_{1-4} alkyl groups; as medicaments.

12. The use of compounds of formula I as claimed in claim 11 as antiinflammatory agents.

13. The use of compounds of formula I as claimed in
25 claim 11 as as anti-allergic agents.

14. The use of compounds of formula I as claimed in claim 11 as immunomodulatory agents.


- 96 -

15. The use of a compound of formula I as claimed in claim 11 in the manufacture of a medicament for use in the treatment of an inflammatory and/or allergic condition and/or diseases with an immunological association.


16. A method of treating inflammatory and/or allergic conditions and/or diseases with an immunological association comprising the administration of a therapeutically effective amount of a compound of formula I as claimed in claim 11 to a mammal in need thereof.

17. A process to prepare a compound of formula I according to claim 1 comprising:

a) reacting a compound of formula IV




15 in which R_1 , R_2 , R_3 and ALK^1 are as previously defined and L_1 represents a leaving group, with a compound of formula V



in which Y, R_4 , ALK^2 , R_5 and R_6 are as previously defined, optionally in the presence of an inert organic liquid, at a temperature in the range 0-250°C, optionally in the presence of a base;


or

b) reacting a compound of formula VI




- in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 and ALK^2 are as initially defined and R_{12}CO represents a group which on reduction yields a group R_5 of formula $-\text{CH}_2\text{R}_{12}$ and R_{11} represents R_6 (or R_{12}CO to give compounds in which R_5 and R_6 are identical), with a reducing agent, at a temperature in the range from -50 to 200°C , to give compounds of formula I in which R_5 represents a group other than hydrogen; or

c) reacting a compound of formula VII




- 10 in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 and ALK^2 are as initially defined and $-\text{N}=\text{R}_{13}$ represents a group which on reduction yields $-\text{NHR}_6$, with a reducing agent, at a temperature in the range 0 – 200°C , to give compounds of formula I in which R_5 represents hydrogen; or
- 15 d) deprotecting a compound of formula VIII




in which R_1 , R_2 , R_3 , ALK^1 , Y , R_4 , R_6 and ALK^2 are as initially defined and PG_1 represents an amine protecting group to give compounds of formula I in which R_5 represents hydrogen; or


- 5 e) reacting a compound of formula IX



in which R_1 , R_2 , R_3 , ALK^1 and Y are as initially defined, with a Wittig reagent, for example (a) a compound of formula Xa




- 10 in which ALK^2 , R_4 , R_5 and R_6 are as initially defined and X represents halo,
or (b) a compound of formula Xb



- 99 -

in which ALK^2 , R_4 , R_5 and R_6 are as initially defined and R_{16} represents an alkyl group, in which Xa or Xb has been pretreated with a base, in an inert organic liquid and the mixture combined in an inert organic liquid, at a temperature in the range from $0^\circ C$ up to the boiling point of the inert organic liquid, to give compounds of formula I in which (CR_4) is connected to Y by a double bond; or


f) reducing compounds of formula XI



10 in which R_1 , R_2 , R_3 , Y, R_4 , ALK^2 , R_5 and R_6 are as initially defined and $-ALK^3-CO-$ represents a group which on reduction yields a group of formula $-ALK^1-$, in an inert organic liquid, at a temperature in the range from $0^\circ C$ up to the boiling point of the inert organic liquid;

15 or


g) hydrolysing compounds of formula XII



in which R_1 , R_2 , R_3 , ALK^1 , R_4 and ALK^2 are as initially defined and PG_2 represents an amine protecting group, optionally in the presence of a base, or an acid to give


compounds of formula I in which R_5 and R_6 represent hydrogen at a temperature in the range 0-250°C; or

h) reducing a compound of formula XIII




5 in which R_1 , R_2 , R_3 , Y , R_4 , R_5 and R_6 are as initially defined and $-(ALK^3)-CO-$ and $(ALK^4)-CO-$ represent groups which on reduction give ALK^1 and ALK^2 respectively, in an inert organic liquid at a temperature in the range from 0°C up to the boiling point of the inert organic liquid; or

10 i) reacting a compound of formula XIV




15 in which R_1 , R_2 , R_3 , ALK^1 are as initially defined, ALK^6 is absent and Y^1 represents a piperidine ring, or a pyridinium ring having a halide counter ion, with a reducing agent, in an inert organic liquid at a temperature in the range from 0°C up to the boiling point of the inert organic liquid to give compounds of formula I in which $---CH_2-$ represents methylene, ALK^2 is absent and R_5 and R_6 represent hydrogen; or

j) reacting a compound of formula XIV




- in which $-(\text{ALK}^6) - \text{CN}$ represents a group which on reduction gives $-\text{CH}(\text{R}_4) - (\text{ALK}^2) - \text{NH}_2$ and Y^1 represents a piperidine ring, or a pyridinium ring having a halide counter ion, with a reducing agent, in an inert organic liquid at a temperature in the range from 0°C up to the boiling point of the inert organic liquid to give compounds of formula I in which the broken line is absent, the free valency on Y is taken up by hydrogen, the free valency on CR_4 is taken up by hydrogen or a C_{1-4} alkyl group, R_5 and R_6 each represent hydrogen and ALK^2 represents a C_{1-4} alkylene chain; or

k) reacting a compound of formula XV




- in which R_1 , R_2 , R_3 , ALK^1 , Y, R_4 and ALK^2 are as initially defined and L_6 represents a leaving group, with a compound of formula XXX




in which R_5 and R_6 are as initially defined, optionally in the presence of an inert organic liquid at a

temperature in the range from 0°C up to the boiling point of the inert organic liquid; or

- 1) reacting a compound of formula XVI




- 5 in which R₁, R₂ and R₃ are as initially defined with a compound of formula XVIII



- 10 in which ALK¹, Y, R₄, ALK², R₅ and R₆ are as initially defined, in the presence of a dialkyl azodicarboxylate, and a phosphorus (III) reagent, optionally in the presence of an inert organic liquid at a temperature in the range from 0°C up to the boiling point of the inert organic liquid; or


- m) reacting compounds of formula XL




- in which R₁, R₂, R₃, ALK¹, Y and R₄ are as initially defined with a reducing agent to give compounds of

formula I in which R_5 and R_6 each represent hydrogen and ALK^2 is absent; or

n) reacting a compound of formula V




5 in which Y, R_4 , ALK^2 , R_5 and R_6 are as initially defined with a compound of formula XLI




10 in which R_1 , R_2 and R_3 are as initially defined and $-(ALK^7)\text{---}CHO$ represents a group which on reaction and reduction yields a group of formula ALK^1 , optionally in the presence of an inert organic liquid at a temperature in the range from 0°C up to the boiling point of the inert organic liquid followed by reaction of the product with a reducing agent, optionally in the presence of an inert organic liquid at a temperature in the range 0-250°C; or

15 o) reacting a compound of formula XVI



in which R_1 , R_2 and R_3 are as initially defined with a compound of formula XLII

- 104 -



in which ALK^1 , Y, R_4 , ALK^2 , R_5 and R_6 are as initially defined and L_8 represents a leaving group, optionally in the presence of an inert organic liquid at a temperature in the range 0-250°C, optionally in the presence of a base; or

- p) alkylating a compound of formula I in which R_6 and/or R_5 , respectively, represents hydrogen either
 - 1) directly using an alkylating agent of formula R_5L_3 or R_6L_3 in which L_3 represents a leaving group in the presence of an organic liquid at a temperature in the range 0-150°C optionally in the presence of a base or
 - 2) by reductive alkylation, at a temperature in the range 0-200°C, optionally in the presence of an inert organic liquid to give compounds of formula I in which R_5 and/or R_6 represents a group other than hydrogen; or
- q) hydrogenating a compound of formula I in which --- represents a double bond, in the presence of an inert organic liquid, at a temperature in the range 0-250°C to give compounds of formula I in which --- represents a single bond.

INTERNATIONAL SEARCH REPORT

 Internat. Application No
 PCT/EP 94/03121

 A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07D211/26 C07D211/70 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

 Minimum documentation searched (classification system followed by classification symbols)
 IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 526 342 (ADIR) 3 February 1993 cited in the application see RN 149069-35-2, 4-Piperidineethanamine, 1-[2-(4-fluorophenoxy)ethyl]- ---	1-17
A	US,A,4 348 401 (BOEHRINGER) 7 September 1982 cited in the application see the whole document ---	1-17
A	US,A,4 027 028 (SCIENCE UNION ET CIE) 31 May 1977 cited in the application see the whole document ---	1-17

-/-

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- '&' document member of the same patent family

Date of the actual completion of the international search

5 January 1995

Date of mailing of the international search report

16. 01. 95

 Name and mailing address of the ISA
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Facs (+ 31-70) 340-3016

Authorized officer

Kissler, B

INTERNATIONAL SEARCH REPORT

Intern. J. Application No.

PCT/EP 94/03121

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 12, no. 426 (C-542) (3273) 10 November 1988 & JP,P,63 154 683 (MITSUBISHI) 27 June 1988 cited in the application see abstract ---	1-17
A	EP,A,0 520 882 (SYNTHELABO) 30 December 1992 cited in the application see RN 147007-82-7, 4-Piperidinemetanamine, 1-[2-(4-fluorophenoxy)ethyl]- ---	1-17
A	GB,A,1 052 302 (THOMAE) 21 December 1966 cited in the application see the whole document -----	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat. Application No
PCT/EP 94/03121

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0526342	03-02-93	FR-A- 2679906 AU-B- 645704 AU-A- 2067692 JP-A- 5208974 NZ-A- 243783 US-A- 5208243	05-02-93 20-01-94 11-03-93 20-08-93 26-07-94 04-05-93
US-A-4348401	07-09-82	DE-A- 2939292 AT-T- 2519 CA-A- 1139759 EP-A, B 0026876 JP-A- 56068669	09-04-81 15-03-83 18-01-83 15-04-81 09-06-81
US-A-4027028	31-05-77	NONE	
EP-A-0520882	30-12-92	FR-A- 2678268 FR-A- 2691148 AU-B- 645697 AU-A- 1858692 AU-B- 645429 AU-A- 1858792 CN-A- 1067887 CN-A- 1067888 EP-A- 0522914 JP-A- 5194419 US-A- 5246939 US-A- 5254560 NZ-A- 243336	31-12-92 19-11-93 20-01-94 14-01-93 13-01-94 07-01-93 13-01-93 13-01-93 13-01-93 03-08-93 21-09-93 19-10-93 25-02-94
GB-A-1052302		CH-A- 436290 CH-A- 436293 FR-M- 3846 FR-A- 1555457 US-A- 3311624	31-01-69